

## When Terrible Things Happen: What You May Experience

### Immediate Reactions

There are a wide variety of positive and negative reactions that survivors can experience during and immediately after a disaster. These include:

| Domain        | Negative Response   | Positive Response   |
|---------------|---|---|
| Cognitive     | Confusion, disorientation, worry, intrusive thoughts and images, self-blame   | Determination and resolve, sharper perception, courage, optimism, faith |
| Emotional     | Shock, sorrow, grief, sadness, fear, anger, numb, irritability, guilt and shame   | Feeling involved, challenged, mobilized                                 |
| Social        | Extreme withdrawal, interpersonal conflict  | Social connectedness, altruistic helping behaviors                      |
| Physiological | Fatigue, headache, muscle tension, stomachache, increased heart rate, exaggerated startle response, difficulties sleeping | Alertness, readiness to respond, increased energy                       |

### Common negative reactions that may continue include:

#### Intrusive reactions

- Distressing thoughts or images of the event while awake or dreaming
- Upsetting emotional or physical reactions to reminders of the experience
- Feeling like the experience is happening all over again (“flashback”)

#### Avoidance and withdrawal reactions

- Avoid talking, thinking, and having feelings about the traumatic event
- Avoid reminders of the event (places and people connected to what happened)
- Restricted emotions; feeling numb
- Feelings of detachment and estrangement from others; social withdrawal
- Loss of interest in usually pleasurable activities

#### Physical arousal reactions

- Constantly being “on the lookout” for danger, startling easily, or being jumpy
- Irritability or outbursts of anger, feeling “on edge”
- Difficulty falling or staying asleep, problems concentrating or paying attention

#### Reactions to trauma and loss reminders

- Reactions to places, people, sights, sounds, smells, and feelings that are reminders of the disaster
- Reminders can bring on distressing mental images, thoughts, and emotional/physical reactions
- Common examples include sudden loud noises, sirens, locations where the disaster occurred, seeing people with disabilities, funerals, anniversaries of the disaster, and television/radio news about the disaster

#### Positive changes in priorities, worldview, and expectations

- Enhanced appreciation that family and friends are precious and important
- Meeting the challenge of addressing difficulties (by taking positive action steps, changing the focus of thoughts, using humor, acceptance)
- Shifting expectations about what to expect from day to day and about what is considered a “good day”
- Shifting priorities to focus more on quality time with family or friends
- Increased commitment to self, family, friends, and spiritual/religious faith

### When a Loved One Dies, Common Reactions Include

- Feeling confused, numb, disbelief, bewildered, or lost
- Feeling angry at the person who died or at people considered responsible for the death
- Strong physical reactions such as nausea, fatigue, shakiness, and muscle weakness
- Feeling guilty for still being alive
- Intense emotions such as extreme sadness, anger, or fear
- Increased risk for physical illness and injury
- Decreased productivity or difficulty making decisions
- Having thoughts about the person who died, even when you don't want to
- Longing, missing, and wanting to search for the person who died
- Children are particularly likely to worry that they or a parent might die
- Children may become anxious when separated from caregivers or other loved ones

### What Helps

- Talking to another person for support or spending time with others
- Engaging in positive distracting activities (sports, hobbies, reading)
- Getting adequate rest and eating healthy meals
- Trying to maintain a normal schedule
- Scheduling pleasant activities
- Taking breaks
- Reminiscing about a loved one who has died
- Focusing on something practical that you can do right now to manage the situation better
- Using relaxation methods (breathing exercises, meditation, calming self-talk, soothing music)
- Participating in a support group
- Exercising in moderation
- Keeping a journal
- Seeking counseling

### What Doesn't Help

- Using alcohol or drugs to cope
- Extreme avoidance of thinking or talking about the event or a death of a loved one
- Violence or conflict
- Overeating or failing to eat
- Excessive TV or computer games
- Blaming others