

# HOCHSCHILD HOMOLOGY RELATIVE TO A FAMILY OF GROUPS

ANDREW NICAS\* AND DAVID ROSENTHAL

ABSTRACT. We define the Hochschild homology groups of a group ring  $\mathbb{Z}G$  relative to a family of subgroups  $\mathcal{F}$  of  $G$ . These groups are the homology groups of a space which can be described as a homotopy colimit, or as a configuration space, or, in the case  $\mathcal{F}$  is the family of finite subgroups of  $G$ , as a space constructed from stratum preserving paths. An explicit calculation is made in the case  $G$  is the infinite dihedral group.

## INTRODUCTION

The *Hochschild homology* of an associative, unital ring  $A$  with coefficients in an  $A$ - $A$  bimodule  $M$  is defined via homological algebra by  $HH_*(A, M) := \mathrm{Tor}_*^{A \otimes A^{\mathrm{op}}}(M, A)$  where  $A^{\mathrm{op}}$  is the opposite ring of  $A$ . In the case  $A = \mathbb{Z}G$ , the integral group ring of a discrete group  $G$ , and  $M = \mathbb{Z}G$ , the Hochschild homology groups  $HH_*(\mathbb{Z}G) := HH_*(\mathbb{Z}G, \mathbb{Z}G)$  have the following homotopy theoretic description. The *cyclic bar construction* associates to a group  $G$  a simplicial set  $N^{\mathrm{cyc}}(G)$  whose homology is  $HH_*(\mathbb{Z}G)$ . Viewing  $G$  as a category,  $\mathbf{G}$ , consisting of a single object and with morphisms identified with the elements of  $G$ , consider the functor  $N$  from  $\mathbf{G}$  to the category of sets given by  $N(*) = G$  and, for a morphism  $g \in G = \mathrm{Mor}_{\mathbf{G}}(*, *)$ , the map  $N(g) : G \rightarrow G$  is conjugation, sending  $x$  to  $g^{-1}xg$ . The geometric realization of  $N^{\mathrm{cyc}}(G)$  is homotopy equivalent to  $\mathrm{hocolim} N$ , the homotopy colimit of  $N$ . There is also a natural homotopy equivalence  $|N^{\mathrm{cyc}}(G)| \rightarrow \mathcal{L}(BG)$  (see [12, Theorem 7.3.11]) where  $BG$  is the classifying space of  $G$  and  $\mathcal{L}(BG)$  is the free loop space of  $BG$ , i.e., the space of continuous maps of the circle into  $BG$ . In particular, there are isomorphisms:

$$HH_*(\mathbb{Z}G) \cong H_*(\mathrm{hocolim} N) \cong H_*(\mathcal{L}(BG)).$$

A *family of subgroups* of a group  $G$  is a non-empty collection of subgroups of  $G$  that is closed under conjugation and finite intersections. In this paper we define the *Hochschild*

---

*Date:* January 31, 2008.

*2000 Mathematics Subject Classification.* 16E40, 19D55, 55R35.

*Key words and phrases.* Hochschild homology, family of subgroups, classifying space.

\*Partially supported by a grant from the Natural Sciences and Engineering Research Council of Canada.

homology of a group ring  $\mathbb{Z}G$  relative to a family of subgroups  $\mathcal{F}$  of  $G$ , denoted  $HH_*^{\mathcal{F}}(\mathbb{Z}G)$ . This is accomplished at the level of spaces. We define a functor  $N_{\mathcal{F}} : \text{Or}(G, \mathcal{F}) \rightarrow \text{CGH}$  where  $\text{Or}(G, \mathcal{F})$  is the *orbit category of  $G$  with respect to  $\mathcal{F}$*  and  $\text{CGH}$  is the category of compactly generated Hausdorff spaces. By definition,  $HH_*^{\mathcal{F}}(\mathbb{Z}G) := H_*(\text{hocolim } N_{\mathcal{F}})$ . If  $\mathcal{F}$  is the trivial family, i.e., contains only the trivial group, then  $N \cong N_{\mathcal{F}}$  and so  $HH_*^{\mathcal{F}}(\mathbb{Z}G) = HH_*(\mathbb{Z}G)$ .

For a discrete group  $G$  and any family  $\mathcal{F}$ , let  $E_{\mathcal{F}}G$  be a universal space for  $G$ -actions with isotropy in  $\mathcal{F}$ . That is,  $E_{\mathcal{F}}G$  is a  $G$ -CW complex whose isotropy groups belong to  $\mathcal{F}$  and for every  $H$  in  $\mathcal{F}$ , the fixed point set  $(E_{\mathcal{F}}G)^H$  is contractible. Given a  $G$ -space  $X$ , let  $F(X)$  be the configuration space of pairs of points in  $X$  which lie on the same  $G$ -orbit. This space inherits a  $G$ -action via restriction of the diagonal action of  $G$  on  $X \times X$ .

Suppose that  $G$  is countable and that the family  $\mathcal{F}$  of subgroups is also countable.

**Theorem A.** *There is a natural homotopy equivalence  $\text{hocolim } N_{\mathcal{F}} \simeq G \backslash F(E_{\mathcal{F}}G)$ .*

Indeed, this homotopy equivalence is a homeomorphism for an appropriate model of the homotopy colimit (see Theorem 3.7 and Corollary 3.8).

Specializing to the case where  $\mathcal{F}$  is the family of finite subgroups of  $G$ , we write  $\underline{E}G := E_{\mathcal{F}}G$  and  $\underline{B}G := G \backslash \underline{E}G$ . Let  $P_{\text{sp}}^{\text{m}}(\underline{B}G)$  denote the space of *marked stratum preserving paths in  $\underline{B}G$*  consisting of stratum preserving paths in  $\underline{B}G$  (with the orbit type partition) whose endpoints are “marked” by an orbit of the diagonal action of  $G$  on  $\underline{E}G \times \underline{E}G$ . We show (see Theorem 4.26(i)):

**Theorem B.** *There is a natural homotopy equivalence  $\text{hocolim } N_{\mathcal{F}} \simeq P_{\text{sp}}^{\text{m}}(\underline{B}G)$ .*

Theorem B is a consequence of Theorem A and a homotopy equivalence  $G \backslash F(X) \simeq P_{\text{sp}}^{\text{m}}(G \backslash X)$  which is valid for any proper  $G$ -CW complex  $X$  (see Theorem 4.20). The Covering Homotopy Theorem of Palais (Theorem 4.7) plays a key role in the proof of the latter result.

If  $\underline{E}G$  satisfies a certain isovariant homotopy theoretic condition then  $P_{\text{sp}}^{\text{m}}(\underline{B}G)$  is homotopy equivalent to a subspace  $\mathcal{L}_{\text{sp}}^{\text{m}}(\underline{B}G) \subset P_{\text{sp}}^{\text{m}}(\underline{B}G)$  which we call the *marked stratified free loop space of  $\underline{B}G$*  (see Theorem 4.26(ii)). We show that this condition is satisfied for appropriate models of  $\underline{E}G$  in the cases:

- (1)  $G$  is torsion free (see Remark 4.25); note that in this case  $\underline{E}G = EG$ , a universal space for free proper  $G$ -actions,

- (2)  $G$  belongs to a particular class of groups which includes the infinite dihedral group and hyperbolic or euclidean triangle groups (see Examples 5.5 and 5.6),
- (3) finite products of such groups (see Remark 5.7).

In the case  $G$  is torsion free,  $\mathcal{L}_{\text{sp}}^{\text{m}}(\underline{B}G)$  is homeomorphic to  $\mathcal{L}(BG)$  by Proposition 4.22 and so our result can be viewed as a generalization of the homotopy equivalence  $|N^{\text{cyc}}(G)| \simeq \mathcal{L}(BG)$ .

There is an equivariant map  $EG \rightarrow \underline{E}G$  which is unique up to equivariant homotopy. This map induces a map  $G \backslash F(EG) \rightarrow G \backslash F(\underline{E}G)$ , equivalently, a map  $\text{hocolim } N \rightarrow \text{hocolim } N_{\mathcal{F}}$ , where  $\mathcal{F}$  is the family of finite subgroups of  $G$ . We explicitly compute this map in the case where  $G = D_{\infty}$ , the infinite dihedral group. In particular, this yields a computation of the homomorphism  $HH_*(\mathbb{Z}D_{\infty}) \rightarrow HH_*^{\mathcal{F}}(\mathbb{Z}D_{\infty})$ ; see Section 6.

The paper is organized as follows. In Section 1 we review some aspects of the theory of homotopy colimits. The functor  $N_{\mathcal{F}} : \text{Or}(G, \mathcal{F}) \rightarrow \text{CGH}$  is defined in Section 2 thus yielding the space  $\mathfrak{N}(G, \mathcal{F}) := \text{hocolim } N_{\mathcal{F}}$  which we call *the Hochschild complex of  $G$  with respect to the family of subgroups  $\mathcal{F}$* . In Section 3 we study the configuration space  $F(X)$  in a general context and give an alternative description of  $\mathfrak{N}(G, \mathcal{F})$  as the orbit space  $G \backslash F(E_{\mathcal{F}}G)$ . The homotopy equivalence  $G \backslash F(X) \simeq P_{\text{sp}}^{\text{m}}(G \backslash X)$ , for any proper  $G$ -CW complex  $X$ , is established in Section 4. We also show in this section that if  $\underline{E}G$  satisfies a certain isovariant homotopy theoretic condition then  $P_{\text{sp}}^{\text{m}}(\underline{B}G)$  is homotopy equivalent to the subspace  $\mathcal{L}_{\text{sp}}^{\text{m}}(\underline{B}G) \subset P_{\text{sp}}^{\text{m}}(\underline{B}G)$ . In Section 5, we show that this condition is satisfied for a class of groups which includes the infinite dihedral group and hyperbolic or euclidean triangle groups. In Section 6 we analyze the map  $G \backslash F(EG) \rightarrow G \backslash F(\underline{E}G)$ , and compute it explicitly in the case  $G = D_{\infty}$  thereby obtaining a computation of the homomorphism  $HH_*(\mathbb{Z}D_{\infty}) \rightarrow HH_*^{\mathcal{F}}(\mathbb{Z}D_{\infty})$ .

## 1. HOMOTOPY COLIMITS AND SPACES OVER A CATEGORY

In this section we provide some categorical preliminaries, following Davis and Lück [7], that will be used in Section 2 to define a Hochschild complex associated to a family of subgroups. Throughout Sections 1 and 2 we work in the category of compactly generated Hausdorff spaces, denoted by  $\text{CGH}$ .<sup>1</sup>

<sup>1</sup>Given a Hausdorff space  $Y$ , *the associated compactly generated space*,  $\text{k}Y$ , is the space with the same underlying set and with the topology defined as follows: a closed set of  $\text{k}Y$  is a set that meets each compact set of  $Y$  in a closed set.  $Y$  is an object of  $\text{CGH}$  if and only if  $Y = \text{k}Y$ , i.e.,  $Y$  is *compactly generated*. The

Let  $\mathcal{C}$  be a small category. A *covariant (contravariant)  $\mathcal{C}$ -space*, is a covariant (contravariant) functor from  $\mathcal{C}$  to CGH. If  $X$  is a contravariant  $\mathcal{C}$ -space and  $Y$  is a covariant  $\mathcal{C}$ -space, then their *tensor product* is defined by

$$X \otimes_{\mathcal{C}} Y = \coprod_{C \in \text{obj}(\mathcal{C})} X(C) \times Y(C) / \sim$$

where  $\sim$  is the equivalence relation generated by

$$(X(\phi)(x), y) \sim (x, Y(\phi)(y))$$

for all  $\phi \in \text{Mor}_{\mathcal{C}}(C, D)$ ,  $x \in X(D)$  and  $y \in Y(C)$ .

A map of  $\mathcal{C}$ -spaces is a natural transformation of functors. Given a  $\mathcal{C}$ -space  $X$  and a topological space  $Z$ , let  $X \times Z$  be the  $\mathcal{C}$ -space defined by  $(X \times Z)(C) = X(C) \times Z$ , where  $C$  is an object in  $\mathcal{C}$ . Two maps of  $\mathcal{C}$ -spaces,  $\alpha, \beta : X \rightarrow X'$ , are  *$\mathcal{C}$ -homotopic* if there is a natural transformation  $H : X \times [0, 1] \rightarrow X'$  such that  $H|_{X \times \{0\}} = \alpha$  and  $H|_{X \times \{1\}} = \beta$ . A map  $\alpha : X \rightarrow X'$  is a  *$\mathcal{C}$ -homotopy equivalence* if there is a map of  $\mathcal{C}$ -spaces  $\beta : X' \rightarrow X$  such that  $\alpha\beta$  is  $\mathcal{C}$ -homotopic to  $\text{id}_{X'}$  and  $\beta\alpha$  is  $\mathcal{C}$ -homotopic to  $\text{id}_X$ . The map  $\alpha : X \rightarrow X'$  is a *weak  $\mathcal{C}$ -homotopy equivalence* if for every object  $C$  in  $\mathcal{C}$ , the map  $\alpha(C) : X(C) \rightarrow X'(C)$  is an ordinary weak homotopy equivalence. Two  $\mathcal{C}$ -spaces  $X$  and  $X'$  are  *$\mathcal{C}$ -homeomorphic* if there are maps  $\alpha : X \rightarrow X'$  and  $\alpha' : X' \rightarrow X$  such that  $\alpha'\alpha = \text{id}_X$  and  $\alpha\alpha' = \text{id}_{X'}$ . If  $X$  and  $X'$  are  $\mathcal{C}$ -homeomorphic contravariant  $\mathcal{C}$ -spaces and  $Y$  and  $Y'$  are  $\mathcal{C}$ -homeomorphic covariant  $\mathcal{C}$ -spaces, then  $X \otimes_{\mathcal{C}} Y$  is homeomorphic to  $X' \otimes_{\mathcal{C}} Y'$ .

A *contravariant free  $\mathcal{C}$ -CW complex*  $X$  is a contravariant  $\mathcal{C}$ -space  $X$  together with a filtration

$$\emptyset = X_{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_n \subset \cdots \subset X = \bigcup_{n \geq 0} X_n$$

such that  $X = \text{colim}_{n \rightarrow \infty} X_n$  and for any  $n \geq 0$ , the  *$n$ -skeleton*,  $X_n$ , is obtained from the  $(n-1)$ -skeleton  $X_{n-1}$  by attaching *free contravariant  $\mathcal{C}$ - $n$ -cells*. That is, there is a

---

product of two spaces  $Y, Z$  in CGH is defined by  $Y \times Z := k(Y \times Z)$  where  $Y \times Z$  on the right side has the product topology. Function space topologies in CGH are defined by applying  $k$  to the compact-open topology. In Sections 3 and 4 we work in the category TOP of all topological spaces and we will have occasion to compare the topologies on  $Y$  and  $kY$  (see Proposition 3.6).

pushout of  $\mathcal{C}$ -spaces of the form

$$\begin{array}{ccc} \coprod_{i \in I_n} \text{Mor}_{\mathcal{C}}(-, C_i) \times S^{n-1} & \longrightarrow & X_{n-1} \\ \downarrow & & \downarrow \\ \coprod_{i \in I_n} \text{Mor}_{\mathcal{C}}(-, C_i) \times D^n & \longrightarrow & X_n \end{array}$$

where  $I_n$  is an indexing set and  $C_i$  is an object in  $\mathcal{C}$ . A *covariant free  $\mathcal{C}$ -CW complex* is defined analogously, the only differences being that the  $\mathcal{C}$ -space is covariant and the  $\mathcal{C}$ -space  $\text{Mor}_{\mathcal{C}}(C_i, -)$  is used in the pushout diagram instead of  $\text{Mor}_{\mathcal{C}}(-, C_i)$ .

A free  $\mathcal{C}$ -CW complex should be thought of as a generalization of a free  $G$ -CW complex. The two notions coincide if  $\mathcal{C}$  is the category associated to the group  $G$ , i.e., the category with one object and one morphism for every element of  $G$ .

Let  $\text{EC}$  be a contravariant free  $\mathcal{C}$ -CW complex such that  $\text{EC}(C)$  is contractible for every object  $C$  of  $\mathcal{C}$ . Such a  $\mathcal{C}$ -space always exists and is unique up to homotopy type [7, Section 3]. One particular example is defined as follows.

Let  $\text{B}^{\text{bar}}\mathcal{C}$  be the *bar construction of the classifying space of  $\mathcal{C}$* , i.e.,  $\text{B}^{\text{bar}}\mathcal{C} = |N\mathcal{C}|$ , the geometric realization of the nerve of  $\mathcal{C}$ . Let  $C$  be an object in  $\mathcal{C}$ . The *undercategory*,  $C \downarrow \mathcal{C}$ , is the category whose objects are pairs  $(f, D)$ , where  $f : C \rightarrow D$  is a morphism in  $\mathcal{C}$ , and whose morphisms,  $p : (f, D) \rightarrow (f', D')$ , consist of a morphism  $p : D \rightarrow D'$  in  $\mathcal{C}$  such that  $p \circ f = f'$ . Notice that a morphism  $\phi : C \rightarrow C'$  induces a functor  $\phi^* : (C' \downarrow \mathcal{C}) \rightarrow (C \downarrow \mathcal{C})$  defined by  $\phi^*(f, D) = (f \circ \phi, D)$ . Let  $\text{E}^{\text{bar}}\mathcal{C} : \mathcal{C} \rightarrow \text{CGH}$  be the contravariant functor defined by

$$\begin{aligned} \text{E}^{\text{bar}}\mathcal{C}(C) &= \text{B}^{\text{bar}}(C \downarrow \mathcal{C}) \\ \text{E}^{\text{bar}}\mathcal{C}(\phi : C \rightarrow C') &= \text{B}^{\text{bar}}(\phi^*) \end{aligned}$$

This is a model for  $\text{EC}$ . Furthermore,  $\text{E}^{\text{bar}}\mathcal{C} \otimes_{\mathcal{C}} *$  is homeomorphic to  $\text{B}^{\text{bar}}\mathcal{C}$  [7, Section 3].

**Lemma 1.1.** [7, Lemma 1.9] *Let  $F : \mathcal{D} \rightarrow \mathcal{C}$  be a covariant functor,  $Z$  a covariant  $\mathcal{D}$ -space and  $X$  a contravariant  $\mathcal{C}$ -space. Let  $F_*Z$  be the covariant  $\mathcal{C}$ -space  $\text{Mor}_{\mathcal{C}}(F(-_{\mathcal{D}}), -_{\mathcal{C}}) \otimes_{\mathcal{D}} Z$ , where  $-_{\mathcal{C}}$  denotes the variable in  $\mathcal{C}$  and  $-_{\mathcal{D}}$  denotes the variable in  $\mathcal{D}$ . Then*

$$X \otimes_{\mathcal{C}} F_*Z \rightarrow (X \circ F) \otimes_{\mathcal{D}} Z$$

*is a homeomorphism.*

*Proof.* The map  $e : X \otimes_{\mathcal{C}} (\text{Mor}_{\mathcal{C}}(F(-\mathcal{D}), -_{\mathcal{C}}) \otimes_{\mathcal{D}} Z) \longrightarrow (X \circ F) \otimes_{\mathcal{D}} Z$  is defined by

$$e([x, [f, y]]) = [X(f)(x), y],$$

where  $x \in X(C)$ ,  $y \in Z(D)$  and  $f \in \text{Mor}_{\mathcal{C}}(F(D), C)$ , for objects  $C$  in  $\mathcal{C}$  and  $D$  in  $\mathcal{D}$ . The inverse is given by mapping  $[w, z] \in (X \circ F) \otimes_{\mathcal{D}} Z$  to  $[w, [\text{id}_{F(D)}, z]]$ , where  $w \in (X \circ F)(D)$  and  $z \in Z(D)$ .  $\square$

**Definition 1.2.** Let  $Y$  be a covariant  $\mathcal{C}$ -space. Then

$$\text{hocolim}_{\mathcal{C}} Y := \text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} Y.$$

A map,  $\alpha : Y \rightarrow Y'$ , of  $\mathcal{C}$ -spaces induces a map  $\alpha_* : \text{hocolim}_{\mathcal{C}} Y \rightarrow \text{hocolim}_{\mathcal{C}} Y'$ . If  $*$  is the  $\mathcal{C}$ -space that sends every object to a point, then

$$\text{hocolim}_{\mathcal{C}} * = \text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} * \cong \text{B}^{\text{bar}} \mathcal{C}.$$

Therefore, the collapse map,  $Y \rightarrow *$ , induces a map  $\bar{\pi} : \text{hocolim}_{\mathcal{C}} Y \rightarrow \text{B}^{\text{bar}} \mathcal{C}$ .

There are several well known constructions for the homotopy colimit, each yielding the same space up to homotopy equivalence (see Talbert [22, Theorem 1.2]). In particular, using the *transport category*,  $\mathcal{T}_{\mathcal{C}}(Y)$ , one can define the homotopy colimit of  $Y$  to be  $\text{B}^{\text{bar}} \mathcal{T}_{\mathcal{C}}(Y)$ . Recall that an object of  $\mathcal{T}_{\mathcal{C}}(Y)$  is a pair  $(C, x)$ , where  $C$  is an object of  $\mathcal{C}$  and  $x \in Y(C)$ , and a morphism  $\phi : (C, x) \rightarrow (C', x')$  is a morphism  $\phi : C \rightarrow C'$  in  $\mathcal{C}$  such that  $Y(\phi)(x) = x'$ . The following lemma shows that  $\text{B}^{\text{bar}} \mathcal{T}_{\mathcal{C}}(Y)$  is not only homotopy equivalent to our definition of the homotopy colimit of  $Y$ , but is in fact homeomorphic to  $\text{hocolim}_{\mathcal{C}} Y$ .

**Lemma 1.3.** *Let  $Y$  be a covariant  $\mathcal{C}$ -space. Then  $\text{E}^{\text{bar}} \mathcal{T}_{\mathcal{C}}(Y) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$  is homeomorphic to  $\text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} Y$ .*

*Proof.* By Lemma 1.1, there is a homeomorphism

$$\text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} \pi_*(*) \rightarrow (\text{E}^{\text{bar}} \mathcal{C} \circ \pi) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$$

where  $\pi : \mathcal{T}_{\mathcal{C}}(Y) \rightarrow \mathcal{C}$  is the projection functor which sends an object  $(C, x)$  to  $C$ . We will show that  $\text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} \pi_*(*)$  is homeomorphic to  $\text{E}^{\text{bar}} \mathcal{C} \otimes_{\mathcal{C}} Y$  and  $(\text{E}^{\text{bar}} \mathcal{C} \circ \pi) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$  is homeomorphic to  $\text{E}^{\text{bar}} \mathcal{T}_{\mathcal{C}}(Y) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$ .

Let  $C$  be an object of  $\mathcal{C}$ . A point in  $\pi_*(*)(C) = \text{Mor}_{\mathcal{C}}(\pi(-), C) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$  is represented by a morphism  $\psi : \pi(D, x) \rightarrow C$  in  $\mathcal{C}$ , where  $(D, x)$  is an object of  $\mathcal{T}_{\mathcal{C}}(Y)$ . Define a natural transformation  $\beta : \pi_*(*) \rightarrow Y$  by  $\beta(C)([\psi]) = Y(\psi)(x)$ . The inverse,  $\beta^{-1} : Y \rightarrow \pi_*(*)$ , is

defined by  $\beta^{-1}(C)(y) = [\text{id}_C]$ , where  $y \in Y(C)$  and  $\text{id}_C : \pi(C, y) \rightarrow C$  is the identity. This induces a homeomorphism  $\text{E}^{\text{bar}}\mathcal{C} \otimes_{\mathcal{C}} \pi_*(*) \rightarrow \text{E}^{\text{bar}}\mathcal{C} \otimes_{\mathcal{C}} Y$ .

Now let  $(C, x)$  be an object of  $\mathcal{T}_{\mathcal{C}}(Y)$ . Then  $(\text{E}^{\text{bar}}\mathcal{C} \circ \pi)(C, x) = \text{E}^{\text{bar}}\mathcal{C}(C) = \text{B}^{\text{bar}}(C \downarrow \mathcal{C})$ , and  $\text{E}^{\text{bar}}\mathcal{T}_{\mathcal{C}}(Y)(C, x) = \text{B}^{\text{bar}}((C, x) \downarrow \mathcal{T}_{\mathcal{C}}(Y))$ . For each  $(C, x)$ , there is an isomorphism of categories  $F_{(C, x)} : C \downarrow \mathcal{C} \rightarrow (C, x) \downarrow \mathcal{T}_{\mathcal{C}}(Y)$  given by  $F_{(C, x)}(f, A) = (f, (A, Y(f)(x)))$ , where  $f : C \rightarrow A$  in  $\mathcal{C}$ . If  $\phi : (f, A) \rightarrow (f', A')$  is a morphism in  $C \downarrow \mathcal{C}$ , then  $F_{(C, x)}(\phi) = \phi : (f, (A, Y(f)(x))) \rightarrow (f', (A', Y(f')(x)))$  is a morphism in  $(C, x) \downarrow \mathcal{T}_{\mathcal{C}}(Y)$  since  $f' = \phi \circ f$ . The inverse of  $F$  is the obvious one. Define the natural transformation  $\alpha : \text{E}^{\text{bar}}\mathcal{C} \circ \pi \rightarrow \text{E}^{\text{bar}}\mathcal{T}_{\mathcal{C}}(Y)$  by  $\alpha(C, x) = \text{B}^{\text{bar}}(F_{(C, x)}) : \text{B}^{\text{bar}}(C \downarrow \mathcal{C}) \rightarrow \text{B}^{\text{bar}}((C, x) \downarrow \mathcal{T}_{\mathcal{C}}(Y))$ , and its inverse by  $\alpha^{-1}(C, x) = \text{B}^{\text{bar}}(F_{(C, x)}^{-1})$ . This induces a homeomorphism  $(\text{E}^{\text{bar}}\mathcal{C} \circ \pi) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} * \rightarrow \text{E}^{\text{bar}}\mathcal{T}_{\mathcal{C}}(Y) \otimes_{\mathcal{T}_{\mathcal{C}}(Y)} *$ .  $\square$

If  $H : \mathcal{D} \rightarrow \mathcal{C}$  is a covariant functor and  $Y$  is a covariant  $\mathcal{C}$ -space, then there is a functor  $\hat{H} : \mathcal{T}_{\mathcal{D}}(Y \circ H) \rightarrow \mathcal{T}_{\mathcal{C}}(Y)$  given by  $\hat{H}(D, x) = (H(D), x)$ . This induces a map  $\text{B}^{\text{bar}}(\hat{H}) : \text{B}^{\text{bar}}\mathcal{T}_{\mathcal{D}}(Y \circ H) \rightarrow \text{B}^{\text{bar}}\mathcal{T}_{\mathcal{C}}(Y)$ . The functor  $H$  also induces a map  $\bar{H} : \text{E}^{\text{bar}}\mathcal{D} \otimes_{\mathcal{D}} Y \circ H \rightarrow \text{E}^{\text{bar}}\mathcal{C} \otimes_{\mathcal{C}} Y$  given by  $\bar{H}([x, y]) = [\text{B}^{\text{bar}}(H_D)(x), y]$ , where  $x \in \text{B}^{\text{bar}}(D \downarrow \mathcal{D})$ ,  $y \in Y(H(D))$  and  $H_D : (D \downarrow \mathcal{D}) \rightarrow (H(D) \downarrow \mathcal{C})$  is the obvious functor induced by  $H$ . The maps  $\text{B}^{\text{bar}}(\hat{H})$  and  $\bar{H}$  are equivalent via the homeomorphism from Lemma 1.3. It is also straightforward to check that the composition of the homeomorphism from Lemma 1.3 with  $\text{B}^{\text{bar}}(\pi) : \text{B}^{\text{bar}}\mathcal{T}_{\mathcal{C}}(Y) \rightarrow \text{B}^{\text{bar}}\mathcal{C}$  is equal to  $\bar{\pi} : \text{hocolim}_{\mathcal{C}} Y \rightarrow \text{B}^{\text{bar}}\mathcal{C}$ .

The transport category definition of the homotopy colimit is employed to prove the following useful lemma.

**Lemma 1.4.** *Let  $H : \mathcal{D} \rightarrow \mathcal{C}$  be a covariant functor and  $Y$  be a covariant  $\mathcal{C}$ -space. Then*

$$(1) \quad \begin{array}{ccc} \text{hocolim}_{\mathcal{D}} Y \circ H & \xrightarrow{\bar{H}} & \text{hocolim}_{\mathcal{C}} Y \\ \pi \downarrow & & \downarrow \bar{\pi} \\ \text{B}^{\text{bar}}\mathcal{D} & \xrightarrow{\text{B}^{\text{bar}}(H)} & \text{B}^{\text{bar}}\mathcal{C} \end{array}$$

*is a pullback diagram.*

*Proof.* Form the pullback diagram

$$\begin{array}{ccc} \mathcal{P}(H, \pi) & \longrightarrow & \mathcal{T}_{\mathcal{C}}(Y) \\ \downarrow & & \downarrow \pi \\ \mathcal{D} & \xrightarrow{H} & \mathcal{C} \end{array}$$

in the category of small categories. The category  $\mathcal{P}(H, \pi)$  is a subcategory of  $\mathcal{T}_{\mathcal{C}}(Y) \times \mathcal{D}$ , where an object  $((C, x), D)$  satisfies  $H(D) = C$ , and a morphism  $(\alpha, \beta) : ((C, x), D) \rightarrow ((C', x'), D')$  satisfies  $\alpha = H(\beta)$ . If  $((C, x), D)$  is an object of  $\mathcal{P}(H, \pi)$ , then  $(C, x)$  is an object of  $\mathcal{T}_{\mathcal{D}}(Y \circ H)$ , and if  $(\alpha, \beta) : ((C, x), D) \rightarrow ((C', x'), D')$  is a morphism of  $\mathcal{P}(H, \pi)$ , then  $\beta : (D, x) \rightarrow (D', x')$  is a morphism of  $\mathcal{T}_{\mathcal{D}}(Y \circ H)$  since  $(Y \circ H)(\beta)(x) = F(\alpha)(x) = x'$ . Hence, we have a functor from  $\mathcal{P}(H, \pi)$  to the transport category  $\mathcal{T}_{\mathcal{D}}(Y \circ H)$  with inverse given by  $(D, x) \mapsto (H(D), x), D$  and  $\beta : (D, x) \rightarrow (D', x') \mapsto (H(\beta), \beta) : (H(D), x), D \rightarrow (H(D'), x'), D'$ . Therefore, we have the pullback diagram

$$\begin{array}{ccc} \mathcal{T}_{\mathcal{D}}(Y \circ H) & \xrightarrow{\hat{H}} & \mathcal{T}_{\mathcal{C}}(Y) \\ \pi \downarrow & & \downarrow \pi \\ \mathcal{D} & \xrightarrow{H} & \mathcal{C} \end{array}$$

Applying  $B^{\text{bar}}$  produces the pullback diagram

$$\begin{array}{ccc} B^{\text{bar}}(\mathcal{T}_{\mathcal{D}}(Y \circ H)) & \longrightarrow & B^{\text{bar}}(\mathcal{T}_{\mathcal{C}}(Y)) \\ \downarrow & & \downarrow \\ B^{\text{bar}}\mathcal{D} & \longrightarrow & B^{\text{bar}}\mathcal{C} \end{array}$$

The result now follows from two applications of Lemma 1.3.  $\square$

## 2. THE ORBIT CATEGORY AND THE HOCHSCHILD COMPLEX

Let  $G$  be a discrete group and  $\mathcal{F}$  a family of subgroups of  $G$  that is closed under conjugation and finite intersections. Let  $\mathcal{O} = \text{Or}(G, \mathcal{F})$  denote the *orbit category of  $G$  with respect to  $\mathcal{F}$* . The objects of  $\mathcal{O}$  are the homogeneous spaces  $G/H$ , with  $H \in \mathcal{F}$ , considered as left  $G$ -sets. Morphisms are all  $G$ -equivariant maps. Therefore,  $\text{Mor}_{\mathcal{O}}(G/H, G/K) = \{r_g \mid g^{-1}Hg \leq K\}$ , where  $r_g$  is right multiplication by  $g$ , i.e.,  $r_g(uH) = (ug)H$  for  $uH \in G/H$ . If  $\mathcal{F}$  is the family of all subgroups of  $G$ , then  $\mathcal{O}$  is called the orbit category. If  $\mathcal{F}$  is taken to be the trivial family, then  $\mathcal{O}$  is the usual category associated to the group  $G$ .

**Definition 2.1** (Hochschild complex of a group associated to a family of subgroups). Let  $\mathcal{O} \times \mathcal{O}$  be the category whose objects are ordered pairs of objects in  $\mathcal{O}$  and whose morphisms are ordered pairs of morphisms in  $\mathcal{O}$ . Let  $\text{Ad} : \mathcal{O} \times \mathcal{O} \rightarrow \text{CGH}$  be the covariant functor defined by

$$\begin{aligned} \text{Ad}(G/H_1, G/H_2) &= H_1 \backslash G/H_2 \\ \text{Ad}(r_{g_1}, r_{g_2})(H_1 u H_2) &= K_1 g_1^{-1} u g_2 K_2, \end{aligned}$$



where  $H_1 \backslash G / H_2$  is the set of  $(H_1, H_2)$  double cosets in  $G$  with the discrete topology and  $(r_{g_1}, r_{g_2}) : (G/H_1, G/H_2) \rightarrow (G/K_1, G/K_2)$  is a morphism in  $\mathcal{O} \times \mathcal{O}$ , and let  $N_{\mathcal{F}} = \text{Ad} \circ \Delta$ , where  $\Delta : \mathcal{O} \rightarrow \mathcal{O} \times \mathcal{O}$  is the diagonal functor. Define

$$\mathfrak{N}(G, \mathcal{F}) = \text{hocolim}_{\mathcal{O}} N_{\mathcal{F}}.$$

We call  $\mathfrak{N}(G, \mathcal{F})$  the *Hochschild complex of  $G$  associated to the family  $\mathcal{F}$* .

*Remark 2.2.* More generally,  $N_{\mathcal{F}}$  can be defined in the case  $G$  is a locally compact topological group and the members of the family of subgroups  $\mathcal{F}$  are closed subgroups of  $G$  by giving  $H_1 \backslash G / H_2$  the quotient topology.

If  $\mathcal{F}$  is the trivial family,  $\{1\}$ , then  $\mathfrak{N}(G, \{1\})$  is homotopy equivalent to  $|N^{\text{cyc}}(G)|$ , the geometric realization of the *cyclic bar construction* ([12, 7.3.10]); indeed, using the 2-sided bar construction as a model for the homotopy colimit of  $N_{\{1\}}$  yields a complex homeomorphic to  $|N^{\text{cyc}}(G)|$ . We refer to  $\mathfrak{N}(G, \{1\})$  as the *classical Hochschild complex of  $G$* .

**Definition 2.3.** The *Hochschild homology of a group ring  $\mathbb{Z}G$  relative to a family of subgroups  $\mathcal{F}$  of  $G$*  is defined to be

$$HH_*^{\mathcal{F}}(\mathbb{Z}G) := H_*(\mathfrak{N}(G, \mathcal{F}); \mathbb{Z}).$$

Using diagram (1) with  $\text{Ad}$  and  $N_{\mathcal{F}}$ , we obtain the following pullback diagram

$$(2) \quad \begin{array}{ccc} \mathfrak{N}(G, \mathcal{F}) & \longrightarrow & \text{hocolim}_{\mathcal{O} \times \mathcal{O}} \text{Ad} \\ \downarrow & & \downarrow \pi \\ \text{B}^{\text{bar}} \mathcal{O} & \xrightarrow{\text{B}^{\text{bar}}(\Delta)} & \text{B}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \end{array}$$

**Lemma 2.4.** Let  $\Delta : \mathcal{O} \rightarrow \mathcal{O} \times \mathcal{O}$  denote the diagonal functor. Then  $\text{hocolim}_{\mathcal{O} \times \mathcal{O}} \text{Ad}$  is homeomorphic to  $(\text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \circ \Delta) \otimes_{\mathcal{O}} *$ .

*Proof.* Let  $T : \mathcal{O} \times \mathcal{O} \rightarrow \text{CGH}$  denote the covariant functor

$$\text{Mor}_{\mathcal{O} \times \mathcal{O}}(\Delta(-\mathcal{O}), -_{\mathcal{O} \times \mathcal{O}}) \otimes_{\mathcal{O}} *.$$

Note that  $\text{Mor}_{\mathcal{O}}(G/L, G/M) = \{r_g \mid g^{-1}Lg \leq M\} \cong \{gM \mid g^{-1}Lg \leq M\}$ . Using this identification, let  $\alpha : \text{Ad} \rightarrow T$  be the natural transformation defined by

$$\alpha(H \backslash G / K)(HgK) = [r_1, r_g],$$

where  $(r_1, r_g) \in \text{Mor}_{\mathcal{O} \times \mathcal{O}}((G/1, G/1), (G/H, G/K))$ . The inverse of  $\alpha$  is given by

$$\alpha^{-1}(G/H, G/K)([r_{g_1}, r_{g_2}]) = Hg_1^{-1}g_2K,$$

where  $(r_{g_1}, r_{g_2}) \in \text{Mor}_{\mathcal{O} \times \mathcal{O}}((G/L, G/L), (G/H, G/K))$  and  $G/L$  is an object in  $\mathcal{O}$ . Thus,  $\text{Ad}$  is naturally equivalent to  $T$ . Therefore,

$$\text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \otimes_{\mathcal{O} \times \mathcal{O}} \text{Ad} \xrightarrow[\cong]{\alpha^*} \text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \otimes_{\mathcal{O} \times \mathcal{O}} T \xrightarrow[\cong]{e} (\text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \circ \Delta) \otimes_{\mathcal{O}} *,$$

where  $e$  is the homeomorphism from Lemma 1.1.  $\square$

**Definition 2.5.** Let  $G$  be a discrete group and  $\mathcal{F}$  be a family of subgroups of  $G$ . A *universal space for  $G$ -actions with isotropy in  $\mathcal{F}$*  is a  $G$ -CW complex,  $\text{E}_{\mathcal{F}}G$ , whose isotropy groups belong to  $\mathcal{F}$  and for every  $H$  in  $\mathcal{F}$ , the fixed point set  $(\text{E}_{\mathcal{F}}G)^H$  is contractible. Such a space is unique up to  $G$ -equivariant homotopy equivalence [14].

Davis and Lück [7, Lemma 7.6] showed that given any model for  $\text{E}\mathcal{O}$ ,  $\text{E}\mathcal{O} \otimes_{\mathcal{O}} \nabla$  is a universal  $G$ -space with isotropy in  $\mathcal{F}$ , where  $\nabla : \mathcal{O} \rightarrow \text{CGH}$  is the covariant functor that sends  $G/H$  to itself and  $r_g : G/H \rightarrow G/K$  to itself.

**Theorem 2.6.** *Let  $G$  be a discrete group and  $\mathcal{F}$  a family of subgroups of  $G$ . Then*

$$\begin{array}{ccc} \mathfrak{N}(G, \mathcal{F}) & \longrightarrow & G \backslash (\mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G) \\ \downarrow & & \downarrow \overline{\rho \times \rho} \\ G \backslash \mathcal{E}_{\mathcal{F}}G & \xrightarrow{\Delta} & G \backslash \mathcal{E}_{\mathcal{F}}G \times G \backslash \mathcal{E}_{\mathcal{F}}G \end{array}$$

is a pullback diagram, where  $\mathcal{E}_{\mathcal{F}}G = \text{E}^{\text{bar}}\mathcal{O} \otimes_{\mathcal{O}} \nabla$ ,  $\rho : \mathcal{E}_{\mathcal{F}}G \rightarrow G \backslash \mathcal{E}_{\mathcal{F}}G$  is the orbit map,  $\overline{\rho \times \rho}$  is the map induced by  $\rho \times \rho$ , and  $\Delta$  is the diagonal map.

*Proof.* There is a homeomorphism,

$$f : (\text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \circ \Delta) \otimes_{\mathcal{O}} * \rightarrow G \backslash (\mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G),$$

defined by  $f([(x, y)]) = q([x, eK], [y, eK])$ , where

$$(x, y) \in \text{B}^{\text{bar}}(G/K \downarrow \mathcal{O}) \times \text{B}^{\text{bar}}(G/K \downarrow \mathcal{O}) \cong (\text{E}^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \circ \Delta)(G/K)$$

and  $q : \mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G \rightarrow G \backslash (\mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G)$  is the orbit map. The inverse of  $f$  is given by  $f^{-1}(q([x, g_1K], [y, g_2K])) = [\text{B}^{\text{bar}}(\epsilon_{g_1K}^*)(x), \text{B}^{\text{bar}}(\epsilon_{g_2K}^*)(y)]$ , where  $\epsilon_{g_iK} : G/1 \rightarrow G/K$

is right multiplication by  $g_i$ . Here we have identified  $B^{\text{bar}}(\mathcal{C} \times \mathcal{D})$  with  $B^{\text{bar}}\mathcal{C} \times B^{\text{bar}}\mathcal{D}$ . Similarly, there is a homeomorphism

$$\bar{f} : B^{\text{bar}}\mathcal{O} \cong E^{\text{bar}}\mathcal{O} \otimes_{\mathcal{O}} * \rightarrow G \backslash \mathcal{E}_{\mathcal{F}}G,$$

defined by  $\bar{f}([x]) = \rho([x, eK])$ , where  $x \in B^{\text{bar}}(G/K \downarrow \mathcal{O})$  and  $\rho : \mathcal{E}_{\mathcal{F}}G \rightarrow G \backslash \mathcal{E}_{\mathcal{F}}G$  is the orbit map. The inverse of  $\bar{f}$  is given by  $(\bar{f})^{-1}(\rho([x, gK])) = [B^{\text{bar}}(\epsilon_{gK}^*)(x)]$ .

Using the homeomorphism from Lemma 2.4, we get the commutative diagram

$$\begin{array}{ccccc} \text{hocolim}_{\mathcal{O} \times \mathcal{O}} \text{Ad} & \xrightarrow[\cong]{e \circ \alpha^*} & (E^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \circ \Delta) \otimes_{\mathcal{O}} * & \xrightarrow[\cong]{f} & G \backslash (\mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G) \\ \downarrow & & & & \downarrow \overline{\rho \times \rho} \\ B^{\text{bar}}(\mathcal{O} \times \mathcal{O}) & \xrightarrow[\cong]{} & B^{\text{bar}}\mathcal{O} \times B^{\text{bar}}\mathcal{O} & \xrightarrow[\bar{f} \times \bar{f}]{\cong} & G \backslash \mathcal{E}_{\mathcal{F}}G \times G \backslash \mathcal{E}_{\mathcal{F}}G \end{array}$$

where  $(\overline{\rho \times \rho})(q(x, y)) = (\rho(x), \rho(y))$ . Since  $B^{\text{bar}}(\Delta)$  composed with the homeomorphism  $B^{\text{bar}}(\mathcal{O} \times \mathcal{O}) \rightarrow B^{\text{bar}}\mathcal{O} \times B^{\text{bar}}\mathcal{O}$  is just the diagonal map  $\Delta : B^{\text{bar}}\mathcal{O} \rightarrow B^{\text{bar}}\mathcal{O} \times B^{\text{bar}}\mathcal{O}$ , diagram (2) completes the proof.  $\square$

*Remark 2.7.* When  $\mathcal{F}$  is the trivial family, the main diagram of Theorem 2.6 becomes:

$$\begin{array}{ccc} \mathfrak{N}(G, \{1\}) & \longrightarrow & G \backslash (EG \times EG) \\ \downarrow & & \downarrow \overline{\rho \times \rho} \\ BG & \xrightarrow{\Delta} & BG \times BG \end{array}$$

Furthermore, in this case, the map  $\overline{\rho \times \rho}$  is a fibration from which it follows that the above square is also a *homotopy* pullback diagram. This observation is part of the folklore of the subject; indeed, one method of establishing the homotopy equivalence  $|N^{\text{cyc}}(G)| \simeq \mathcal{L}(BG)$  involves replacing  $\overline{\rho \times \rho}$  with the fibration  $BG^I \rightarrow BG \times BG$  given by evaluation at endpoints where  $BG^I$  is the space of paths in  $BG$ . For a general family  $\mathcal{F}$ , Theorem 2.6 is, to our knowledge, new and we note that the map  $\overline{\rho \times \rho}$  in Theorem 2.6 is typically not a fibration.

If  $\mathcal{F}' \subset \mathcal{F}$ , then there is an inclusion functor  $\iota : \text{Or}(G, \mathcal{F}') \rightarrow \text{Or}(G, \mathcal{F})$ . Clearly,  $N_{\mathcal{F}'} = N_{\mathcal{F}} \circ \iota$ , which induces a map  $\mathfrak{N}(G, \mathcal{F}') \rightarrow \mathfrak{N}(G, \mathcal{F})$ . This map is examined in Section 6 in the case when  $\mathcal{F}'$  is the trivial family and  $\mathcal{F}$  is the family of finite subgroups.

### 3. THE CONFIGURATION SPACE $F(X)$

In this section we investigate, in a general context, some basic properties of the configuration space,  $F(X)$ , of pairs of points in a  $G$ -space  $X$  which lie on the same  $G$ -orbit.

Let  $G$  be a topological group. The *category of left  $G$ -spaces*, denoted by  ${}_G\text{TOP}$ , is the category whose objects are left  $G$ -spaces, i.e., topological spaces  $X$  together with a continuous left  $G$ -action  $G \times X \rightarrow X$ , written as  $(g, x) \mapsto gx$ , and whose morphisms are continuous equivariant maps  $f : X \rightarrow Y$ . Henceforth, we abbreviate “left  $G$ -space” to “ $G$ -space”.

Given a  $G$ -space  $X$ , define  $A_X : G \times X \rightarrow X \times X$  by  $A_X(g, x) := (x, gx)$  for  $(g, x) \in G \times X$ . Note that  $A_X$  is continuous and  $G$ -equivariant where  $G \times X$  is given the left  $G$  action

$$(3) \quad h(g, x) := (hgh^{-1}, hx) \text{ for } h, g \in G \text{ and } x \in X$$

and  $X \times X$  is given the diagonal  $G$ -action. Hence the image of  $A_X$  is a  $G$ -invariant subspace of  $X \times X$ .

**Definition 3.1.** Define  $F : {}_G\text{TOP} \rightarrow {}_G\text{TOP}$  on an object  $X$  by  $F(X) := \text{image}(A_X)$  with the left  $G$ -action inherited from diagonal  $G$ -action on  $X \times X$ . If  $f : X \rightarrow Y$  is equivariant, i.e., a morphism in  ${}_G\text{TOP}$ , then the diagram

$$\begin{array}{ccc} G \times X & \xrightarrow{A_X} & X \times X \\ \text{id}_G \times f \downarrow & & \downarrow f \times f \\ G \times Y & \xrightarrow{A_Y} & Y \times Y \end{array}$$

is commutative and so  $f \times f$  restricts to an equivariant map  $F(f) : F(X) \rightarrow F(Y)$ . Clearly,  $F(\text{id}_X) = \text{id}_{F(X)}$  and  $F(f_1 f_2) = F(f_1) F(f_2)$  for composable morphisms  $f_1$  and  $f_2$ , i.e.,  $F$  is a functor.

Note that  $F(X)$  is the subspace of  $X \times X$  consisting of those pairs  $(x, y)$  such that  $x$  and  $y$  lie in the same orbit of the  $G$ -action.

There is an evident natural isomorphism  $F(X) \times I \cong F(X \times I)$ , where  $I$  is the unit interval with the trivial  $G$ -action, given by  $((x, y), t) \mapsto ((x, t), (y, t))$  for  $(x, y) \in F(X)$  and  $t \in I$ . If  $H : X \times I \rightarrow Y$  is an equivariant homotopy then

$$F(X) \times I \xrightarrow{\cong} F(X \times I) \xrightarrow{F(H)} Y$$

is an equivariant homotopy from  $F(H_0)$  to  $F(H_1)$ , where  $H_t := H(-, t)$ . Hence  $F$  factors through the homotopy category of  $G$  TOP with the following consequence.

**Proposition 3.2.** *If the map  $f : X \rightarrow Y$  is an equivariant homotopy equivalence then  $F(f) : F(X) \rightarrow F(Y)$  is an equivariant homotopy equivalence.*  $\square$

**Definition 3.3.** In the category TOP of all topological spaces we use the following notation for the *standard pullback construction*. Given maps  $e : A \rightarrow Z$  and  $f : B \rightarrow Z$  define  $E(e, f) := \{(x, y) \in A \times B \mid e(x) = f(y)\}$  topologized as a subspace of  $A \times B$  with the product topology. The maps  $p_1 : E(e, f) \rightarrow A$  and  $p_2 : E(e, f) \rightarrow B$  are given, respectively, by the restriction of the projections  $A \times B \rightarrow A$  and  $A \times B \rightarrow B$ . The square

$$\begin{array}{ccc} E(e, f) & \xrightarrow{p_2} & B \\ p_1 \downarrow & & \downarrow f \\ A & \xrightarrow{e} & Z \end{array}$$

is a pullback diagram in TOP which we refer to as a *standard pullback diagram*.

**Proposition 3.4.** *There is a pullback diagram*

$$\begin{array}{ccc} F(X) & \xrightarrow{i} & X \times X \\ q \downarrow & & \downarrow \rho \times \rho \\ G \backslash X & \xrightarrow{\Delta} & G \backslash X \times G \backslash X \end{array}$$

where  $i$  is the inclusion  $F(X) = \text{image}(A_X) \subset X \times X$ ,  $\rho : X \rightarrow G \backslash X$  is the orbit map,  $\Delta$  is the diagonal map and  $q : F(X) \rightarrow G \backslash X$  is given by  $q((x, y)) = \rho(y)$  for  $(x, y) \in F(X)$ .

*Proof.* The standard pullback construction yields

$$E(\Delta, \rho \times \rho) = \{(\rho(x), x_1, x_2) \in (G \backslash X) \times X \times X \mid \rho(x) = \rho(x_1) = \rho(x_2)\}.$$

The map  $j : F(X) \rightarrow E(\Delta, \rho \times \rho)$  given by  $j((x, y)) = (\rho(x), x, y)$  is a homeomorphism with inverse  $(\rho(x), x, y) \mapsto (x, y)$ . Also  $p_1 j = q$  and  $p_2 j = i$  where  $p_1 : E(\Delta, \rho \times \rho) \rightarrow G \backslash X$  and  $p_2 : E(\Delta, \rho \times \rho) \rightarrow X \times X$  are the restrictions of the corresponding projections.  $\square$

The space  $G \backslash F(X)$  can also be described as a pullback as follows:

**Theorem 3.5.** *There is a pullback diagram*

$$\begin{array}{ccc} G \backslash F(X) & \xrightarrow{\bar{i}} & G \backslash (X \times X) \\ \bar{q} \downarrow & & \downarrow \overline{\rho \times \rho} \\ G \backslash X & \xrightarrow{\Delta} & G \backslash X \times G \backslash X \end{array}$$

where  $\bar{i}$ ,  $\bar{q}$  and  $\overline{\rho \times \rho}$  are induced by  $i$ ,  $q$  and  $\rho \times \rho$  respectively (as in Proposition 3.4).

*Proof.* The pullback diagram of Proposition 3.4 factors as:

$$\begin{array}{ccc} F(X) & \xrightarrow{i} & X \times X \\ q' \downarrow & & \downarrow \rho' \\ E(\Delta, \overline{\rho \times \rho}) & \xrightarrow{p_2} & G \backslash (X \times X) \\ p_1 \downarrow & & \downarrow \overline{\rho \times \rho} \\ G \backslash X & \xrightarrow{\Delta} & G \backslash X \times G \backslash X \end{array}$$

where  $\rho' : X \times X \rightarrow G \backslash (X \times X)$  is the orbit map,  $q'((x, y)) = (\rho(x), \rho'(x, y))$  for  $(x, y) \in F(X)$  and  $E(\Delta, \overline{\rho \times \rho})$  together with the maps  $p_1, p_2$  is the standard pullback construction. The outer square in the above diagram is a pullback by Proposition 3.4 and the lower square is a pullback by construction. It follows that the upper square is a pullback. By Lemma 3.18,  $q'$  induces a homeomorphism  $G \backslash F(X) \cong E(\Delta, \overline{\rho \times \rho})$ .  $\square$

A Hausdorff space  $X$  is *compactly generated* if a set  $A \subset X$  is closed if and only if it meets each compact set of  $X$  in a closed set.

**Proposition 3.6.** *Suppose that  $G$  is a countable discrete group and that  $X$  is a countable  $G$ -CW complex i.e.,  $X$  has countably many  $G$ -cells. Then  $F(X)$  and  $G \backslash F(X)$  are compactly generated Hausdorff spaces.*

*Proof.* Milnor showed that the product of two countable CW complexes is a CW complex, [18, Lemma 2.1]. Since  $X$  and  $G \backslash X$  are countable CW complexes,  $G \backslash X \times X \times X$  is also a CW complex and thus compactly generated. By Proposition 3.4,  $F(X)$  is homeomorphic to a closed subset of this space and hence must be compactly generated. The space  $X \times X$  is a CW complex and so  $G \backslash (X \times X)$  is also a CW complex because the diagonal  $G$ -action on  $X \times X$  is cellular. By Theorem 3.5,  $G \backslash F(X)$  is homeomorphic to a closed subset of the CW complex  $G \backslash X \times G \backslash (X \times X)$  and hence must be compactly generated.  $\square$

Recall that for a discrete group  $G$  and family of subgroups  $\mathcal{F}$ , we denote the bar construction model for the universal space for  $G$ -actions with isotropy in  $\mathcal{F}$  by  $\mathcal{E}_{\mathcal{F}}G$  (see Theorem 2.6).

**Theorem 3.7.** *Suppose that  $G$  is a countable discrete group and that  $\mathcal{F}$  is a countable family of subgroups. Then there is a natural homeomorphism  $\mathfrak{N}(G, \mathcal{F}) \cong G \backslash F(\mathcal{E}_{\mathcal{F}}G)$ .*

*Proof.* By Theorem 3.5, there is a pullback diagram in TOP:

$$\begin{array}{ccc} G \backslash F(\mathcal{E}_{\mathcal{F}}G) & \longrightarrow & G \backslash (\mathcal{E}_{\mathcal{F}}G \times \mathcal{E}_{\mathcal{F}}G) \\ \downarrow & & \downarrow \overline{\rho \times \rho} \\ G \backslash \mathcal{E}_{\mathcal{F}}G & \xrightarrow{\Delta} & G \backslash \mathcal{E}_{\mathcal{F}}G \times G \backslash \mathcal{E}_{\mathcal{F}}G \end{array}$$

Since  $G$  and  $\mathcal{F}$  are countable,  $\mathcal{E}_{\mathcal{F}}G$  is a countable CW complex. All the spaces appearing in the above diagram are compactly generated by Proposition 3.6 and its proof. It follows that this diagram is also a pullback diagram in the category of compactly generated Hausdorff spaces. A comparison with the pullback diagram in the statement of Theorem 2.6 yields a natural homeomorphism  $\mathfrak{N}(G, \mathcal{F}) \cong G \backslash F(\mathcal{E}_{\mathcal{F}}G)$ .  $\square$

**Corollary 3.8.** *Suppose that  $G$  is a countable discrete group and that  $\mathcal{F}$  is a countable family of subgroups. Let  $E_{\mathcal{F}}G$  be any  $G$ -CW model for the universal space for  $G$ -actions with isotropy in  $\mathcal{F}$ . Then there is a natural homotopy equivalence  $\mathfrak{N}(G, \mathcal{F}) \simeq G \backslash F(E_{\mathcal{F}}G)$ .*

*Proof.* There is an equivariant homotopy equivalence  $J : \mathcal{E}_{\mathcal{F}}G \rightarrow E_{\mathcal{F}}G$ , which is unique up to equivariant homotopy. By Proposition 3.2,  $J$  induces a homotopy equivalence  $G \backslash F(\mathcal{E}_{\mathcal{F}}G) \rightarrow G \backslash F(E_{\mathcal{F}}G)$ . Composition with the homeomorphism of Theorem 3.7 yields the conclusion.  $\square$

Note that in Corollary 3.8, “natural” means that for an inclusion  $\mathcal{F}' \subset \mathcal{F}$  of families of subgroups of  $G$ , the corresponding square diagram is homotopy commutative.

Recall that a continuous map  $f : Y \rightarrow Z$  is *proper* if for any topological space  $W$   $f \times \text{id}_W : Y \times W \rightarrow Z \times W$  is a closed map (equivalently,  $f$  is a closed map with quasi-compact fibers, [4, I, 10.2, Theorem 1(b)]). There are several distinct notions of a “proper action” of a topological group on a topological space; see [2] for their comparison. We will use the following definition ([4, III, 4.1, Definition 1]).

**Definition 3.9.** A left action of a topological group  $G$  on a topological space  $X$  is *proper* provided the map  $A_X : G \times X \rightarrow X \times X$  is proper in which case we say that  $X$  is a *proper  $G$ -space*.

**Proposition 3.10.** *Suppose that the topological group  $G$  acts freely and properly on the  $G$ -space  $X$ . Then  $A_X : G \times X \rightarrow F(X)$  is a homeomorphism. Consequently,  $A_X$  induces a homeomorphism  $\bar{A}_X : G \backslash (G \times X) \rightarrow G \backslash F(X)$  where the  $G$ -action on  $G \times X$  is given by (3).*

*Proof.* Clearly,  $A_X$  is a continuous surjection. Since the  $G$ -action is proper,  $A_X$  is a closed map. If  $A_X(g_1, x_1) = A_X(g_2, x_2)$  then  $x_1 = x_2$  and  $g_1x_1 = g_2x_2$ . Since the  $G$ -action is free,  $g_1 = g_2$  and so  $A_X$  is injective. It follows that  $A_X$  is a homeomorphism.  $\square$

Let  $\text{conj}(G)$  denote the set of conjugacy classes of the group  $G$ . For  $g \in G$ , let  $C(g) \in \text{conj}(G)$  denote the conjugacy class of  $g$  and let  $Z(g) := \{h \in G \mid hg = gh\}$  denote the centralizer of  $g$ .

**Proposition 3.11.** *Suppose that  $G$  is a discrete group acting on a topological space  $X$ . Then there is a homeomorphism*

$$G \backslash (G \times X) \cong \coprod_{C(g) \in \text{conj}(G)} Z(g) \backslash X$$

where the right side of the isomorphism is a disjoint topological sum.

*Proof.* The space  $G \times X$  is the disjoint union of the  $G$ -invariant subspaces  $C(g) \times X$ ,  $C(g) \in \text{conj}(G)$ . Since  $G$  is discrete,  $C(g) \times X$  is both open and closed in  $G \times X$ . It follows that  $G \backslash (G \times X)$  is the disjoint topological sum of the spaces  $G \backslash (C(g) \times X)$ ,  $C(g) \in \text{conj}(G)$ . The map  $G \backslash (C(g) \times X) \rightarrow Z(g) \backslash X$  which takes the  $G$ -orbit of  $(hgh^{-1}, x)$  to the  $Z(g)$ -orbit of  $h^{-1}x$  is a homeomorphism whose inverse is the map which takes the  $Z(g)$ -orbit of  $x \in X$  to the  $G$ -orbit of  $(g, x)$ .  $\square$

Combining Propositions 3.10 and 3.11 yields:

**Corollary 3.12.** *Suppose that  $G$  is a discrete group which acts freely and properly on a topological space  $X$ . Then there is a homeomorphism*

$$G \backslash F(X) \cong \coprod_{C(g) \in \text{conj}(G)} Z(g) \backslash X$$

where the right side of the isomorphism is a disjoint topological sum.  $\square$



*Remark 3.13.* A discrete group  $G$  acts freely and properly on a space  $X$  if and only if  $G \backslash X$  is Hausdorff and the orbit map  $\rho : X \rightarrow G \backslash X$  is a covering projection.

As a consequence of Corollary 3.12, if a non-trivial discrete group  $G$  acts freely and properly on a non-empty topological space  $X$  then  $G \backslash F(X)$  is never connected. However, if  $G$  acts properly but *not* freely then  $F(X)$ , hence also  $G \backslash F(X)$ , can be connected; see Examples 5.5 and 5.6.

**Definition 3.14.** Let  $X$  be a  $G$ -space. The subspace  $F(X)_0 \subset F(X)$  is defined to be the union of the connected components of  $F(X)$  which meet the *diagonal* of  $X \times X$ , i.e., the subspace  $\Delta(X) = \{(x, x) \in X \times X\}$ . In particular, if  $X$  is connected then  $F(X)_0$  is the connected component of  $F(X)$  containing  $\Delta(X)$ .

**Proposition 3.15.**  $F(X)_0$  is a  $G$ -invariant subspace of  $F(X)$ .

*Proof.* Let  $C$  be a component of  $F(X)$  such that  $C \cap \Delta(X) \neq \emptyset$ . Left translation by  $g \in G$ ,  $L_g : F(X) \rightarrow F(X)$ , is a homeomorphism and so  $L_g(C)$  is also a component of  $F(X)$ .  $\emptyset \neq L_g(C \cap \Delta(X)) = L_g(C) \cap \Delta(X)$  and so  $L_g(C) \subset F(X)_0$ .  $\square$

*Remark 3.16.* Suppose that the discrete group  $G$  acts freely and properly on  $X$ . Then by Proposition 3.10, the map  $A_X : G \times X \rightarrow F(X)$  is an equivariant homeomorphism and  $F(X)_0 = A_X(\{1\} \times X) = \Delta(X)$ .

The remainder of this section is devoted to the proof of various elementary lemmas which have been employed above.

**Lemma 3.17.** Consider the standard pullback diagram:

$$\begin{array}{ccc} E(f, p) & \xrightarrow{p_2} & Y \\ p_1 \downarrow & & \downarrow p \\ Z & \xrightarrow{f} & X \end{array}$$

If  $p$  is an open map then  $p_1$  is also an open map.

*Proof.* Let  $V \subset X$  and  $W \subset Y$  be open sets. Then  $p_1((V \times W) \cap E(f, p)) = V \cap f^{-1}(p(W))$ . Note that  $f^{-1}(p(W))$  is open since the map  $p$  is open and  $f$  is continuous and so  $V \cap f^{-1}(p(W))$  is also open. Since sets of the form  $(V \times W) \cap E(f, p)$  give a basis for the topology of  $E(f, p)$  and  $p_1$  preserves unions, the conclusion follows.  $\square$

**Lemma 3.18.** *Let  $G$  be a topological group, let  $Y$  be a  $G$ -space and let  $f : Z \rightarrow G \backslash Y$  be a continuous map. Consider the standard pullback diagram:*

$$\begin{array}{ccc} E(f, \rho) & \xrightarrow{p_2} & Y \\ p_1 \downarrow & & \downarrow \rho \\ Z & \xrightarrow{f} & G \backslash Y \end{array}$$

where  $\rho : Y \rightarrow G \backslash Y$  is the orbit map and  $G$  acts on  $E(f, \rho)$  by  $g(z, y) := (z, gy)$  for  $g \in G$  and  $(z, y) \in E(f, \rho)$ . Then  $p_1$  induces a homeomorphism  $\bar{p}_1 : G \backslash E(f, \rho) \rightarrow Z$  given by  $\bar{p}_1(q(z, y)) = z$  for  $(z, y) \in E(f, \rho)$  where  $q : E(f, \rho) \rightarrow G \backslash E(f, \rho)$  is the orbit map.

*Proof.* The map  $\bar{p}_1$  is clearly well defined and continuous since  $p_1 = \bar{p}_1 q$  and  $G \backslash E(f, \rho)$  has the identification topology determined by the orbit map  $q$ . Since  $\rho$  is surjective,  $p_1$  is surjective and thus  $\bar{p}_1$  is also surjective. Suppose  $\bar{p}_1(q(z_1, x_1)) = \bar{p}_1(q(z_2, x_2))$ . Then  $z_1 = z_2$  and so  $\rho(x_1) = f(z_1) = f(z_2) = \rho(x_2)$ . Hence  $q(z_1, x_2) = q(z_2, x_2)$ , demonstrating that  $\bar{p}_1$  is injective. Since  $\rho$  is an open map,  $p_1$  is also an open map by Lemma 3.17. Let  $U \subset G \backslash E(f, \rho)$  be open. Since  $q$  is surjective,  $U = q(q^{-1}(U))$ . Hence

$$\bar{p}_1(U) = \bar{p}_1 q(q^{-1}(U)) = p_1(q^{-1}(U))$$

which is open since  $q^{-1}(U)$  is open and  $p_1$  is an open map. Hence  $\bar{p}_1$  is an open map. It follows that  $\bar{p}_1$  is a homeomorphism.  $\square$

#### 4. THE MARKED STRATIFIED FREE LOOP SPACE

Suppose that  $X$  is a proper  $G$ -CW complex where  $G$  is a discrete group. In this section, we show that the orbit space  $G \backslash F(X)$  is homotopy equivalent to the space,  $P_{\text{sp}}^m(G \backslash X)$ , of stratum preserving paths in  $G \backslash X$  whose endpoints are “marked” by an orbit of the diagonal action of  $G$  on  $X \times X$  (see Theorem 4.20). The Covering Homotopy Theorem of Palais plays a key role in the proof of this result. If  $X$  satisfies a suitable isovariant homotopy theoretic condition then  $P_{\text{sp}}^m(G \backslash X)$  is shown to be homotopy equivalent to a subspace  $\mathcal{L}_{\text{sp}}^m(G \backslash X) \subset P_{\text{sp}}^m(G \backslash X)$  which we call the *marked stratified free loop space of  $G \backslash X$*  (see Theorem 4.23). Applying these results to the case  $X = \underline{E}G$ , a universal space for proper  $G$ -actions, yields a homotopy equivalence between the homotopy colimit,  $\mathfrak{N}(G, \mathcal{F})$ , of Section 2 and  $P_{\text{sp}}^m(G \backslash \underline{E}G)$  and also, for suitable  $G$ , to  $\mathcal{L}_{\text{sp}}^m(G \backslash X)$  (see Theorem 4.26).

**4.1. Orbit maps as stratified fibrations.** We recall some of the basic definitions from the theory of stratified spaces following the treatment in [11].

**Definition 4.1.** A *partition* of a topological space  $X$  consists of an indexing set  $\mathcal{J}$  and a collection  $\{X_j \mid j \in \mathcal{J}\}$  of pairwise disjoint subspaces of  $X$  such that  $X = \bigcup_{j \in \mathcal{J}} X_j$ . For each  $j \in \mathcal{J}$ ,  $X_j$  is called the  *$j$ -th stratum*.

A *refinement* of a partition  $\{X_j \mid j \in \mathcal{J}\}$  of a space  $X$  is another partition  $\{X'_i \mid i \in \mathcal{J}'\}$  of  $X$  such that for every  $i \in \mathcal{J}'$  there exists  $j \in \mathcal{J}$  such that  $X'_i \subset X_j$ . The *component refinement* of a partition  $\{X_j \mid j \in \mathcal{J}\}$  of  $X$  is the refinement obtained by taking the  $X'_i$ 's to be the connected components of the  $X_j$ 's.

**Definition 4.2.** A *stratification* of a topological space  $X$  is a locally finite partition  $\{X_j \mid j \in \mathcal{J}\}$  of  $X$  such that each  $X_j$  is locally closed in  $X$ . We say that  $X$  together with its stratification is a *stratified space*.

If  $X$  is a space with a given partition then a map  $f : Z \times A \rightarrow X$  is *stratum preserving along  $A$*  if for each  $z \in Z$ ,  $f(\{z\} \times A)$  lies in a single stratum of  $X$ . In particular, a map  $f : Z \times I \rightarrow X$  is a *stratum preserving homotopy* if it is stratum preserving along  $I$ .

A *class of topological spaces* will mean a subclass of the class of all topological spaces, typically defined by a property, for example, the class of all metrizable spaces.

**Definition 4.3.** Let  $X$  and  $Y$  be spaces with given partitions. A map  $p : X \rightarrow Y$  is a *stratified fibration* with respect to a class of topological spaces  $\mathcal{W}$  if for any space  $Z$  in  $\mathcal{W}$  and any commutative square

$$\begin{array}{ccc} Z & \xrightarrow{f} & X \\ i_0 \downarrow & & \downarrow p \\ Z \times I & \xrightarrow{H} & Y \end{array}$$

where  $i_0(z) := (z, 0)$  and  $H$  is a stratum preserving homotopy, there exists a stratum preserving homotopy  $\tilde{H} : Z \times I \rightarrow X$  such that  $\tilde{H}(z, 0) = f(z)$  for all  $z \in Z$  and  $p\tilde{H} = H$ .

**Definition 4.4.** Let  $X$  be a space with a given partition. The *space of stratum preserving paths in  $X$* , denoted by  $P_{\text{sp}}(X)$ , is the subspace of  $X^I$ , the space of continuous maps of the unit interval into  $X$  with the compact-open topology, consisting of stratum preserving paths, i.e., paths  $\omega : I \rightarrow X$  such that  $\omega(I)$  belongs to a single stratum of  $X$ .

Observe that a homotopy  $H : Z \times I \rightarrow X$  is stratum preserving if and only if its adjoint  $\widehat{H} : Z \rightarrow X^I$ , given by  $\widehat{H}(z)(t) := H(z, t)$  for  $(z, t) \in Z \times I$ , has  $\widehat{H}(Z) \subset P_{\text{sp}}(X)$ .

A group action on a space determines an invariant partition on that space as follows.

**Definition 4.5** (Orbit type partition). Let  $G$  be a topological group and let  $X$  be a  $G$ -space. For a subgroup  $H \subset G$ , let  $X_H := \{x \in X \mid G_x = H\}$  where  $G_x$  is the isotropy subgroup at  $x$ . Let  $(H) := \{gHg^{-1} \mid g \in G\}$ , the set of conjugates of  $H$  in  $G$ , and  $X_{(H)} := \bigcup_{K \in (H)} X_K$ . Let  $\mathcal{J}$  denote the set of conjugacy classes of subgroups of  $G$  of the form  $(G_x)$ . The subspaces  $X_{(H)}$  are  $G$ -invariant and  $\{X_{(H)} \mid (H) \in \mathcal{J}\}$  is a partition of  $X$  called the *orbit type partition* of  $X$ . Let  $\rho : X \rightarrow G \backslash X$  denote the orbit map. The set  $\{\rho(X_{(H)}) \mid (H) \in \mathcal{J}\}$  is a partition of  $G \backslash X$  also called the *orbit type partition* of  $G \backslash X$ .

*Remark 4.6.* If  $G$  is a Lie group acting smoothly and properly on a smooth manifold  $M$  then the component refinement of the orbit type partition of  $M$  is a stratification of  $M$  which, in addition, satisfies Whitney's Conditions A and B; see [8, Theorem 2.7.4].

An equivariant map  $f : X \rightarrow Y$  between two  $G$ -spaces is *isovariant* if for every  $x \in X$ ,  $G_x = G_{f(x)}$ . An equivariant homotopy  $H : X \times I \rightarrow Y$  is said to be *isovariant* if for each  $t \in I$ ,  $H_t := H(-, t)$  is isovariant.

We make use of the following version of the Covering Homotopy Theorem of Palais.

**Theorem 4.7** (Covering Homotopy Theorem). *Let  $G$  be a Lie group, let  $X$  be a  $G$ -space and let  $Y$  be a proper  $G$ -space. Assume that every open subset of  $G \backslash X$  is paracompact. Suppose that  $f : X \rightarrow Y$  is an isovariant map and that  $F : G \backslash X \times I \rightarrow G \backslash Y$  is a homotopy such that  $F_0 \circ \rho_X = \rho_Y \circ f$ , where  $\rho_X : X \rightarrow G \backslash X$  and  $\rho_Y : Y \rightarrow G \backslash Y$  are the orbit maps, and  $F(\rho_X(X_{(H)}) \times I) \subset \rho_Y(Y_{(H)})$  for every compact subgroup  $H \subset G$ . Then there exists an isovariant homotopy  $\widetilde{F} : X \times I \rightarrow Y$  such that  $\widetilde{F}_0 = f$  and  $F \circ (\rho_X \times \text{id}_I) = \rho_Y \circ \widetilde{F}$ .  $\square$*

*Remark 4.8.* The Covering Homotopy Theorem (CHT) was originally demonstrated by Palais in the case  $G$  is a compact Lie group and  $X$  and  $Y$  are second countable and locally compact ([19, 2.4.1]). Palais later observed ([20, 4.5]) that his proof of the CHT generalizes to the case of proper actions of a non-compact Lie group. Bredon proved the CHT under the hypotheses that  $G$  is compact and that  $G \backslash X$  has the property that every open subset is paracompact ([5, II, Theorem 7.3]). A topological space is *hereditarily paracompact* if every subspace is paracompact, equivalently, if every *open* subspace is paracompact ([16, App. I,

Lemma 8]). The class of hereditarily paracompact spaces includes all metric spaces (since any metric space is paracompact) and all CW complexes ([16, II, sec. 4]). The authors of [1] observed that Bredon's proof of [5, II, Theorem 7.1], from which the CHT is deduced, can be adapted to the case of a proper action of a non-compact Lie group; see the discussion following [1, Theorem 1.5]. Also, note that it is not necessary to assume that the  $G$ -action on  $X$  is proper because the induced  $G$ -action on the standard pullback  $E(F, \rho_Y)$  is proper by Lemma 4.9 below.

**Lemma 4.9.** *Suppose that  $G \times Y \rightarrow Y$  is a proper action of a topological group  $G$  on a Hausdorff space  $Y$ . Let  $Z$  be a Hausdorff space and  $f : Z \rightarrow G \backslash Y$  a continuous map. Let  $\rho : Y \rightarrow G \backslash Y$  denote the orbit map. Then the induced action of  $G$  on the standard pullback  $E(f, \rho)$  is proper.*

*Proof.* By hypothesis, the map  $A_Y : G \times Y \rightarrow Y \times Y$ ,  $A_Y(g, y) = (y, gy)$ , is proper. Since  $Z$  is Hausdorff, the diagonal map  $\Delta : Z \rightarrow Z \times Z$  is proper. The product of two proper maps is proper and thus  $A_Y \times \Delta : G \times Y \times Z \rightarrow Y \times Y \times Z \times Z$  is proper. It follows that  $A_{Z \times Y} = h_2 \circ (A_Y \times \text{id}_Z) \circ h_1 : G \times Z \times Y \rightarrow Z \times Y \times Z \times Y$  is proper where  $h_1 : G \times Z \times Y \rightarrow G \times Y \times Z$  and  $h_2 : Y \times Y \times Z \times Z \rightarrow Z \times Y \times Z \times Y$  are the “interchange” homeomorphisms,  $h_1(g, z, y) = (g, y, z)$  and  $h_2(y_1, y_2, z_1, z_2) = (z_1, y_1, z_2, y_2)$ . Since the action of  $G$  on  $Y$  is proper,  $G \backslash Y$  is Hausdorff ([4, III, 4.2, Proposition 3]) and so  $E(f, \rho)$  is a closed subset of  $Z \times Y$ . Hence the restriction of  $A_{Z \times Y}$  to  $G \times E(f, \rho)$  is a proper map. This restriction map factors as  $i \circ A_{E(f, \rho)}$  where  $i : E(f, \rho) \times E(f, \rho) \hookrightarrow Z \times Y \times Z \times Y$  is inclusion and thus  $A_{E(f, \rho)}$  is a proper map ([4, I, 10.2, Proposition 5(d)]).  $\square$

**Theorem 4.10.** *Suppose that  $G$  is a Lie group and that  $Y$  is a proper  $G$ -space. Let  $Y$  and  $G \backslash Y$  have the orbit type partitions. Then the orbit map  $\rho : Y \rightarrow G \backslash Y$  is a stratified fibration with respect to the class of hereditarily paracompact spaces.*

*Proof.* Let  $Z$  be a hereditarily paracompact space, let  $F : Z \times I \rightarrow G \backslash Y$  be a homotopy which is stratum preserving along  $I$  and let  $f : Z \rightarrow Y$  be a map such that  $\rho \circ f = F_0$ . Consider the standard pullback diagram:

$$\begin{array}{ccc} E(F_0, \rho) & \xrightarrow{p_2} & Y \\ p_1 \downarrow & & \downarrow \rho \\ Z & \xrightarrow{F_0} & G \backslash Y \end{array}$$

By Lemma 3.18,  $p_1$  induces a homeomorphism  $\bar{p}_1 : G \backslash E(F_0, \rho) \rightarrow Z$ . The map  $p_2$  is clearly isovariant. The CHT (Theorem 4.7) implies that there is an isovariant homotopy  $\tilde{F} : E(F_0, \rho) \times I \rightarrow Y$  such that  $\rho \circ \tilde{F} = F \circ (p_1 \times \text{id}_I)$  and  $\tilde{F}_0 = p_2$ . Define  $\hat{f} : Z \rightarrow E(F_0, \rho)$  by  $\hat{f}(z) = (z, f(z))$  for  $z \in Z$ . Let  $\bar{F} : Z \times I \rightarrow Y$  be given by  $\bar{F} = \tilde{F} \circ (\hat{f} \times \text{id}_I)$ . Then  $\rho \circ \bar{F} = F$  and  $\bar{F}_0 = f$ ; furthermore,  $\bar{F}$  is stratum preserving along  $I$ .  $\square$

**Corollary 4.11.** *Suppose that  $G$  is a Lie group and that  $Y$  is a proper  $G$ -space. Let  $H \subset G$  be a subgroup. Then the orbit map  $\rho : Y_{(H)} \rightarrow G \backslash Y_{(H)}$  is a Serre fibration.*

*Proof.* Suppose that  $Z$  is a compact polyhedron. Then  $Z$  is metrizable and thus hereditarily paracompact. Given a homotopy  $F : Z \times I \rightarrow G \backslash Y_{(H)}$  and a map  $f : Z \rightarrow Y_{(H)}$  such that  $F_0 = \rho \circ f$ , apply Theorem 4.10 to  $j \circ F$  and  $i \circ f$ , where  $i : Y_{(H)} \hookrightarrow Y$  and  $j : G \backslash Y_{(H)} \hookrightarrow G \backslash Y$  are the inclusions, to obtain  $\tilde{F} : Z \times I \rightarrow Y_{(H)}$  with  $\rho \circ \tilde{F} = F$  and  $\tilde{F}_0 = f$ .  $\square$

**4.2. Spaces of marked stratum preserving paths.** We apply the results of Section 4.1 in the case  $G$  is a discrete group to show that, for a proper  $G$ -CW complex  $X$ , the orbit space  $G \backslash F(X)$  is homotopy equivalent to the space,  $P_{\text{sp}}^{\text{m}}(G \backslash X)$ , of stratum preserving paths in  $G \backslash X$  whose endpoints are “marked” by an orbit of the diagonal action of  $G$  on  $X \times X$ ; see Theorem 4.20. That theorem together with Corollaries 3.8 and 4.24 are used to prove Theorem 4.26, which subsumes Theorem B as stated in the introduction to this paper.

**Lemma 4.12.** *Suppose that  $G$  is a discrete group and that  $Y$  is a proper  $G$ -space. Then the orbit map  $\rho : Y \rightarrow G \backslash Y$  has the unique path lifting property for stratum preserving paths, i.e., given a stratum preserving path  $\omega : I \rightarrow G \backslash Y$  and  $y \in \rho^{-1}(\omega(0))$  there exists a unique path  $\tilde{\omega} : I \rightarrow Y$  such that  $\tilde{\omega}(0) = y$  and  $\rho \circ \tilde{\omega} = \omega$ .*

*Proof.* Let  $\omega : I \rightarrow G \backslash Y$  be a stratum preserving path, i.e., there exists a finite subgroup  $H \subset G$  such that  $\omega(I) \subset \rho(Y_{(H)}) = G \backslash Y_{(H)}$ . By Corollary 4.11, the restriction of  $\rho$  to  $Y_{(H)}$ ,  $\rho : Y_{(H)} \rightarrow G \backslash Y_{(H)}$ , is a Serre fibration. The fiber over  $\rho(y)$ , where  $y \in Y_{(H)}$ , is the orbit  $G \cdot y$  which is discrete since the  $G$ -action on  $Y$  is proper. By [21, 2.2 Theorem 5], a fibration with discrete fibers has the unique path lifting property (note that in the cited theorem, the given fibration is assumed to be a Hurewicz fibration; however, the proof of this theorem uses only the homotopy lifting property respect to  $I$  and so remains valid for a Serre fibration).  $\square$

Combining Theorem 4.10 and Lemma 4.12 yields:

**Proposition 4.13.** (Unique Lifting.) *Suppose that  $G$  is a discrete group and that  $Y$  is a proper  $G$ -space. Let  $Z$  be a hereditarily paracompact space. Suppose that  $F : Z \times I \rightarrow G \backslash Y$  is stratum preserving homotopy and that  $f : Z \rightarrow Y$  is a map such that  $\rho \circ f = F_0$ . Then there exists a unique stratum preserving homotopy  $\tilde{F} : Z \times I \rightarrow Y$  such that  $\rho \circ \tilde{F} = F$  and  $\tilde{F}_0 = f$ .  $\square$*

We define a “stratified homotopy” version of  $F(X)$  as follows.

**Definition 4.14.** Let  $X$  be a  $G$ -space with its orbit type partition. The  $G$ -space  $F_{\text{sp}}(X)$  is given by:

$$F_{\text{sp}}(X) := \{(\omega, y) \in P_{\text{sp}}(X) \times X \mid \text{there exists } g \in G \text{ such that } y = g\omega(1)\}$$

where  $G$  acts on  $F_{\text{sp}}(X)$  by the restriction of the diagonal action of  $G$  on  $P_{\text{sp}}(X) \times X$ .

Note that there is a pullback diagram

$$\begin{array}{ccc} F_{\text{sp}}(X) & \xrightarrow{i} & P_{\text{sp}}(X) \times X \\ q \downarrow & & \downarrow (\rho \circ \text{ev}_1) \times \rho \\ G \backslash X & \xrightarrow{\Delta} & G \backslash X \times G \backslash X \end{array}$$

where  $i$  is the inclusion  $F_{\text{sp}}(X) \hookrightarrow P_{\text{sp}}(X) \times X$ ,  $\rho : X \rightarrow G \backslash X$  is the orbit map,  $\text{ev}_1 : P_{\text{sp}}(X) \rightarrow X$  is evaluation at 1,  $\Delta$  is the diagonal map and  $q : F_{\text{sp}}(X) \rightarrow G \backslash X$  is given by  $q((\omega, y)) = \rho(y)$  for  $(\omega, y) \in F_{\text{sp}}(X)$ .

**Proposition 4.15.** *The map  $\ell : F(X) \rightarrow F_{\text{sp}}(X)$  given by  $\ell(x, y) = (c_x, y)$ , where  $c_x$  is the constant path at  $x$ , is an equivariant homotopy equivalence with an equivariant homotopy inverse  $j : F_{\text{sp}}(X) \rightarrow F(X)$  given by  $j(\omega, y) = (\omega(1), y)$ .*

*Proof.* Observe that  $j \circ \ell = \text{id}_{F(X)}$ . Define a homotopy,  $H : F_{\text{sp}}(X) \times I \rightarrow F_{\text{sp}}(X)$  by  $H((\omega, y), t) = (\omega_t, y)$  where  $\omega_t \in P_{\text{sp}}(X)$  is the path  $\omega_t(s) = \omega((1-s)t + s)$  for  $s \in I$ . Then  $H$  is an equivariant homotopy from  $\text{id}_{F_{\text{sp}}(X)}$  to  $\ell \circ j$ .  $\square$

**Corollary 4.16.** *The map  $\ell : F(X) \rightarrow F_{\text{sp}}(X)$  induces a homotopy equivalence  $\bar{\ell} : G \backslash F(X) \rightarrow G \backslash F_{\text{sp}}(X)$ .  $\square$*

If  $G$  is a Lie group, we say that a  $G$ -CW complex  $X$  is *proper* if  $G$  acts properly on  $X$ . By [13, Theorem 1.23], a  $G$ -CW complex  $X$  is proper if and only if for each  $x$  in  $X$  the isotropy group  $G_x$  is compact. In particular, if  $G$  is discrete, then  $X$  is a proper  $G$ -CW complex if and only if  $G_x$  is finite for every  $x$  in  $X$ .

**Proposition 4.17.** *Let  $G$  be a discrete group. Suppose that  $X$  is a proper  $G$ -CW complex. Then there is a pullback diagram:*

$$\begin{array}{ccc} F_{\text{sp}}(X) & \xrightarrow{q_2} & X \times X \\ q_1 \downarrow & & \downarrow \rho \times \rho \\ P_{\text{sp}}(G \backslash X) & \xrightarrow{\text{ev}_{0,1}} & G \backslash X \times G \backslash X \end{array}$$

where  $\rho : X \rightarrow G \backslash X$  is the orbit map,  $q_1$  and  $q_2$  are given, respectively, by  $q_1(\omega, y) = \rho \circ \omega$  and  $q_2(\omega, y) = (\omega(0), y)$  for  $(\omega, y) \in F_{\text{sp}}(X)$ , and  $\text{ev}_{0,1}(\tau) = (\tau(0), \tau(1))$  for  $\tau \in P_{\text{sp}}(G \backslash X)$ .

*Proof.* Let  $Z$  be a hereditarily paracompact space. Suppose  $h = (h_0, h_1) : Z \rightarrow X \times X$  and  $f : Z \rightarrow P_{\text{sp}}(G \backslash X)$  are maps such that  $\text{ev}_{0,1} f = (\rho \times \rho)h$ . Let  $\check{f} : Z \times I \rightarrow G \backslash X$  be the adjoint of  $f$ , i.e.,  $\check{f}(z, t) = f(z)(t)$  for  $(z, t) \in Z \times I$ . Note that  $\check{f}$  is stratum preserving along  $I$ . The diagram

$$\begin{array}{ccc} Z & \xrightarrow{h_0} & X \\ i_0 \downarrow & & \downarrow \rho \\ Z \times I & \xrightarrow{\check{f}} & G \backslash X \end{array}$$

is commutative where  $i_0(z) = (z, 0)$  for  $z \in Z$ . By Proposition 4.13, there exists a unique  $F : Z \times I \rightarrow X$  which is stratum preserving along  $I$  such that  $\rho F = \check{f}$  and  $F i_0 = h_0$ . Let  $\widehat{F} : Z \rightarrow P_{\text{sp}}(X)$  be the adjoint of  $F$ . Then  $Q : Z \rightarrow F_{\text{sp}}(X)$  given by  $Q(z) = (\widehat{F}(z), h_1(z))$  for  $z \in Z$  is the unique map such that  $h = q_2 Q$  and  $f = q_1 Q$ . In order to conclude that the diagram appearing in the statement of the proposition is a pullback diagram in TOP, it suffices to show that the spaces  $F_{\text{sp}}(X)$  and

$$E(\text{ev}_{0,1}, \rho \times \rho) = \{(\omega, x, y) \in P_{\text{sp}}(G \backslash X) \times X \times X \mid \omega(0) = \rho(x), \omega(1) = \rho(y)\}$$

are hereditarily paracompact. Since  $X$  and  $G \backslash X$  are CW complexes, the main theorem of [6] implies that the path spaces  $X^I$  and  $(G \backslash X)^I$  are *stratifiable* in the sense of [3, Definition 1.1] (despite the sound alike terminology, this notion of “stratifiable” is not directly related to our Definition 4.2). It is shown in [3] that any CW complex is stratifiable, that a countable product of stratifiable spaces is stratifiable and that a stratifiable space



is paracompact and *perfectly normal*, i.e., normal and every closed set is a countable intersection of open sets. Hence  $X^I \times X$  and  $(G \backslash X)^I \times X \times X$  are stratifiable and thus paracompact and perfectly normal. A subspace of a paracompact and perfectly normal space is also paracompact and perfectly normal ([16, App. I, Theorem 10]). In particular,  $F_{\text{sp}}(X) \subset X^I \times X$  and  $E(\text{ev}_{0,1}, \rho \times \rho) \subset (G \backslash X)^I \times X \times X$  and all of their subspaces are paracompact.  $\square$

**Definition 4.18.** The space  $P_{\text{sp}}^{\text{m}}(G \backslash X)$  of *marked stratum preserving paths in  $G \backslash X$*  consists of stratum preserving paths in  $G \backslash X$  whose endpoints are “marked” by an orbit of the diagonal action of  $G$  on  $X \times X$ . More precisely,  $P_{\text{sp}}^{\text{m}}(G \backslash X) = E(\text{ev}_{0,1}, \overline{\rho \times \rho})$  where

$$\begin{array}{ccc} E(\text{ev}_{0,1}, \overline{\rho \times \rho}) & \xrightarrow{p_2} & G \backslash (X \times X) \\ p_1 \downarrow & & \downarrow \overline{\rho \times \rho} \\ P_{\text{sp}}(G \backslash X) & \xrightarrow{\text{ev}_{0,1}} & G \backslash X \times G \backslash X \end{array}$$

is a standard pullback diagram and  $\overline{\rho \times \rho}$  is induced by  $\rho \times \rho : X \times X \rightarrow G \backslash X \times G \backslash X$ .

**Proposition 4.19.** *Let  $G$  be a discrete group. Suppose that  $X$  is a proper  $G$ -CW complex. Then the map  $q : F_{\text{sp}}(X) \rightarrow P_{\text{sp}}^{\text{m}}(G \backslash X)$  given by  $q(\omega, y) = (\rho \circ \omega, \rho'(\omega(0), y))$ , where  $\rho' : X \times X \rightarrow G \backslash (X \times X)$  is the orbit map of the diagonal action, induces a homeomorphism  $\bar{q} : G \backslash F_{\text{sp}}(X) \rightarrow P_{\text{sp}}^{\text{m}}(G \backslash X)$ .*

*Proof.* The pullback diagram of Proposition 4.17 factors as:

$$\begin{array}{ccc} F_{\text{sp}}(X) & \xrightarrow{q_2} & X \times X \\ q \downarrow & & \downarrow \rho' \\ P_{\text{sp}}^{\text{m}}(G \backslash X) & \xrightarrow{p_2} & G \backslash (X \times X) \\ p_1 \downarrow & & \downarrow \overline{\rho \times \rho} \\ P_{\text{sp}}(G \backslash X) & \xrightarrow{\text{ev}_{0,1}} & G \backslash X \times G \backslash X \end{array}$$

The outer square in the above diagram is a pullback by Proposition 4.17 and the lower square is a pullback by definition. It follows that the upper square is a pullback. By Lemma 3.18,  $q$  induces a homeomorphism  $\bar{q} : G \backslash F_{\text{sp}}(X) \rightarrow P_{\text{sp}}^{\text{m}}(G \backslash X)$ .  $\square$

Combining Corollary 4.16 and Proposition 4.19 yields:

**Theorem 4.20.** *Let  $G$  be a discrete group. Suppose that  $X$  is a proper  $G$ -CW complex. Then the map  $\bar{q} \circ \bar{\ell} : G \backslash F(X) \rightarrow P_{\text{sp}}^{\text{m}}(G \backslash X)$  is a homotopy equivalence.*  $\square$

**Definition 4.21.** The *stratified free loop space* of  $G \backslash X$ , denoted by  $\mathcal{L}_{\text{sp}}(G \backslash X)$ , is the subspace of  $P_{\text{sp}}(G \backslash X)$  consisting of closed paths, i.e.,  $\omega \in P_{\text{sp}}(G \backslash X)$  such that  $\omega(0) = \omega(1)$ . The *marked stratified free loop space* of  $G \backslash X$ , denoted by  $\mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X)$ , is the subspace of  $P_{\text{sp}}^{\text{m}}(G \backslash X)$  given by:

$$\mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X) = \{(\omega, \rho'(x, y)) \in P_{\text{sp}}^{\text{m}}(G \backslash X) \mid (x, y) \in F(X)_0\}.$$

(Recall that  $\rho : X \rightarrow G \backslash X$  and  $\rho' : X \times X \rightarrow G \backslash (X \times X)$  are the orbit maps and that  $F(X)_0$  is the union of the components of  $F(X)$  meeting the diagonal.) Note that if  $(\omega, \rho'(x, y)) \in \mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X)$  then  $\omega(0) = \rho(x) = \rho(y) = \omega(1)$  and so  $\omega \in \mathcal{L}_{\text{sp}}(G \backslash X)$ .

There is a standard pullback diagram:

$$\begin{array}{ccc} \mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X) & \xrightarrow{p_2} & G \backslash F(X)_0 \\ p_1 \downarrow & & \downarrow p \\ \mathcal{L}_{\text{sp}}(G \backslash X) & \xrightarrow{\text{ev}_0} & G \backslash X \end{array}$$

where  $p$  is given by  $p(\rho'(x, y)) = \rho(x)$  for  $\rho'(x, y) \in G \backslash F(X)_0$ .

Let  $\bar{\Delta} : G \backslash X \rightarrow G \backslash (X \times X)$  denote the map induced by the diagonal map,  $\Delta : X \rightarrow X \times X$ . Define the map  $\iota : \mathcal{L}_{\text{sp}}(G \backslash X) \rightarrow \mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X)$  by  $\iota(\omega) = (\omega, \bar{\Delta}(\omega(0)))$ . The composite  $p_1 \iota$  is the identity map of  $\mathcal{L}_{\text{sp}}(G \backslash X)$  and so  $\mathcal{L}_{\text{sp}}(G \backslash X)$  is homeomorphic to a retract of  $\mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X)$ . In general,  $\iota$  is not a homotopy equivalence; for example, in the case of the infinite dihedral group,  $D_{\infty}$ , acting on  $\mathbb{R}$  as in Example 5.5,  $\mathcal{L}_{\text{sp}}(D_{\infty} \backslash \mathbb{R})$  is contractible whereas  $\mathcal{L}_{\text{sp}}^{\text{m}}(D_{\infty} \backslash \mathbb{R})$  is not simply connected.

**Proposition 4.22.** *If the discrete group  $G$  acts freely and properly on  $X$  then the map  $\iota : \mathcal{L}_{\text{sp}}(G \backslash X) \rightarrow \mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X)$  is a homeomorphism; furthermore,  $\mathcal{L}_{\text{sp}}(G \backslash X) = \mathcal{L}(G \backslash X)$ , the space of closed paths in  $G \backslash X$ .*

*Proof.* Since the  $G$ -action on  $X$  is free and proper, by Remark 3.16,  $F(X)_0$  is the diagonal of  $X \times X$  and so  $p : G \backslash F(X)_0 \rightarrow G \backslash X$  is a homeomorphism. Thus  $p_1 : \mathcal{L}_{\text{sp}}^{\text{m}}(G \backslash X) \rightarrow \mathcal{L}_{\text{sp}}(G \backslash X)$  is also homeomorphism since it is a pullback of  $p$ . Hence  $\iota = (p_1)^{-1}$  is a homeomorphism. Since the  $G$ -action is free, there is only one stratum and so  $\mathcal{L}_{\text{sp}}(G \backslash X) = \mathcal{L}(G \backslash X)$ .  $\square$

Define  $\tilde{S}$  to be the image of the map  $G \times P_{\text{sp}}(X) \rightarrow X \times X$  given by  $(g, \sigma) \mapsto (\sigma(0), g\sigma(1))$ . Note that  $\tilde{S}$  is a  $G$ -invariant subset of  $X \times X$  and that  $F(X) \subset \tilde{S}$ .

**Theorem 4.23.** *Suppose that the pair  $(\tilde{S}, F(X)_0)$  can be deformed isovariantly into the pair  $(F(X)_0, F(X)_0)$ , i.e., there is an isovariant homotopy  $H : \tilde{S} \times I \rightarrow \tilde{S}$  such that*

$H(-, 0)$  is the identity of  $\tilde{S}$  and  $H(\tilde{S} \times \{1\} \cup F(X)_0 \times I) \subset F(X)_0$ . Then the inclusion  $i : \mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X) \hookrightarrow P_{\text{sp}}^{\text{m}}(G \setminus X)$  is a homotopy equivalence.

*Proof.* Let  $H : \tilde{S} \times I \rightarrow \tilde{S}$  be an isovariant homotopy such that  $H(-, 0)$  is the identity of  $\tilde{S}$  and  $H(\tilde{S} \times \{1\} \cup F(X)_0 \times I) \subset F(X)_0$ . Write  $H = (H_1, H_2)$  where  $H_j : \tilde{S} \times I \rightarrow X$  for  $j = 1, 2$ . Define the homotopy  $b : P_{\text{sp}}^{\text{m}}(G \setminus X) \times I \rightarrow P_{\text{sp}}^{\text{m}}(G \setminus X)$  by

$$b((\omega, \rho'(x, y)), s)(t) = \begin{cases} \rho \circ H_1((x, y), s - 3t) & \text{if } 0 \leq t \leq s/3, \\ \omega(\frac{3t-s}{3-2s}) & \text{if } s/3 \leq t \leq 1 - s/3, \\ \rho \circ H_2((x, y), s + 3t - 3) & \text{if } 1 - s/3 \leq t \leq 1. \end{cases}$$

where  $\rho : X \rightarrow G \setminus X$  and  $\rho' : X \times X \rightarrow G \setminus (X \times X)$  are the orbit maps. Define the homotopy  $B : P_{\text{sp}}^{\text{m}}(G \setminus X) \times I \rightarrow P_{\text{sp}}^{\text{m}}(G \setminus X)$  by

$$B((\omega, \rho'(x, y)), s) = (b((\omega, \rho'(x, y)), s), \rho'(H((x, y), s))).$$

The hypotheses on  $H$  imply that  $B$  is a deformation of the pair  $(P_{\text{sp}}^{\text{m}}(G \setminus X), \mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X))$  into the pair  $(\mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X), \mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X))$  and so  $i : \mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X) \hookrightarrow P_{\text{sp}}^{\text{m}}(G \setminus X)$  is a homotopy equivalence.  $\square$

The inclusion  $F(X)_0 \hookrightarrow \tilde{S}$  is an *isovariant strong deformation retract* if there is a homotopy  $H : \tilde{S} \times I \rightarrow \tilde{S}$  as in Theorem 4.23 with the additional property that  $H$  is stationary along  $F(X)_0$ .

**Corollary 4.24.** *If  $F(X)_0 \hookrightarrow \tilde{S}$  is an isovariant strong deformation retract then  $i : \mathcal{L}_{\text{sp}}^{\text{m}}(G \setminus X) \hookrightarrow P_{\text{sp}}^{\text{m}}(G \setminus X)$  is a homotopy equivalence.*  $\square$

*Remark 4.25.* Suppose in Theorem 4.23 that the discrete group  $G$  acts freely and properly. Then  $\tilde{S} = X \times X$  and  $F(X)_0 = \Delta(X)$ , the diagonal of  $X \times X$ ; see Remark 3.16. The hypothesis of Theorem 4.23 asserts that  $(X \times X, \Delta(X))$  is deformable into  $(\Delta(X), \Delta(X))$  and so the diagonal map  $\Delta : X \rightarrow X \times X$  is a homotopy equivalence. This implies that  $X$  is contractible and hence a model for the universal space,  $EG$ , for free  $G$ -actions, provided  $X$  has the equivariant homotopy type of a  $G$ -CW complex. Conversely, suppose that  $EG$  is a  $G$ -CW model for the universal space such that  $EG \times EG$  with the product topology and the diagonal  $G$ -action is also a  $G$ -CW complex and has an equivariant subdivision such that  $\Delta(EG)$  is a subcomplex. Then  $\Delta(EG) \subset EG \times EG$  is an equivariant, hence isovariant (since the  $G$ -action is free), strong deformation retract.

In Section 5 we show that the hypothesis of Corollary 4.24 is satisfied for a class of groups which includes the infinite dihedral group and hyperbolic or euclidean triangle groups and where  $X$  is a universal space for  $G$ -actions with finite isotropy.

**Theorem 4.26.** *Suppose that  $G$  is a countable discrete group and that  $\mathcal{F}$  is its family of finite subgroups. Let  $\underline{EG} := E_{\mathcal{F}}G$ , a universal space for proper  $G$ -actions, and  $\underline{BG} := G \backslash \underline{EG}$ .*

- (i) *There is a homotopy equivalence  $\mathfrak{N}(G, \mathcal{F}) \simeq P_{\text{sp}}^{\text{m}}(\underline{BG})$ .*
- (ii) *If  $\underline{EG}$  satisfies the hypothesis of Corollary 4.24 then there is a homotopy equivalence  $\mathfrak{N}(G, \mathcal{F}) \simeq \mathcal{L}_{\text{sp}}^{\text{m}}(\underline{BG})$ .*

*Proof.* Conclusion (i) of the theorem is a direct consequence of Corollary 3.8 and Theorem 4.20. Conclusion (ii) follows from (i) and Corollary 4.24.  $\square$

If  $G$  is torsion free then the family  $\mathcal{F}$  of finite subgroups of  $G$  is the trivial family and so  $|N^{\text{cyc}}(G)| \simeq \mathfrak{N}(G, \mathcal{F})$  and  $\mathcal{L}_{\text{sp}}^{\text{m}}(\underline{BG}) \cong \mathcal{L}(\underline{BG})$  (Proposition 4.22); furthermore, by Remark 4.25, Theorem 4.26(ii) applies thus recovering the familiar result  $|N^{\text{cyc}}(G)| \simeq \mathcal{L}(\underline{BG})$ .

## 5. EXAMPLES

Let  $\underline{EG}$  denote the universal space for proper  $G$ -actions and  $\underline{BG} = G \backslash \underline{EG}$ . In this section, we show that if  $G$  is the infinite dihedral group or a hyperbolic or euclidean triangle group, then the hypothesis of Corollary 4.24 is satisfied; that is,  $F(\underline{EG})_0 \hookrightarrow \tilde{S}$  is an isovariant strong deformation retract. By Theorem 4.26, this implies that  $\mathfrak{N}(G, \mathcal{F}) \simeq P_{\text{sp}}^{\text{m}}(\underline{BG}) \simeq \mathcal{L}_{\text{sp}}^{\text{m}}(\underline{BG})$ , where  $\mathcal{F}$  is the family of finite subgroups of  $G$ . This is accomplished by showing that, for these groups,  $F(\underline{EG})$  is path connected and  $F(X) \hookrightarrow \tilde{S}$  is a  $G \times G$ -isovariant strong deformation retract.

Let  $G$  be a discrete group and  $X$  a proper  $G$ -space. Recall that  $F(X)$  is the image of  $A_X : G \times X \rightarrow X \times X$ , where  $A_X(g, x) := (x, gx)$  for  $(g, x) \in G \times X$ , and  $\tilde{S}$  is the image of the map  $G \times P_{\text{sp}}(X) \rightarrow X \times X$  given by  $(g, \sigma) \mapsto (\sigma(0), g\sigma(1))$ . Notice that  $F(X)$  and  $\tilde{S}$  are each  $G \times G$ -invariant subsets of  $X \times X$ . Let  $\rho : X \rightarrow G \backslash X$  denote the orbit map. Then  $F(X) = (\rho \times \rho)^{-1}(\Delta(G \backslash X))$ , and  $\tilde{S} = (\rho \times \rho)^{-1}(\{(\sigma(0), \sigma(1)) \mid \sigma \in P_{\text{sp}}(G \backslash X)\})$  by Lemma 4.12.

**Proposition 5.1.** *Let  $G$  be a discrete group and  $X$  a proper  $G$ -space. Assume that  $G \backslash X$  is homeomorphic to a subset of  $\mathbb{R}^n$  for some  $n$ , and that the images of the strata of  $G \backslash X$  in  $\mathbb{R}^n$  are convex. Then  $F(X) \hookrightarrow \tilde{S}$  is a  $G \times G$ -isovariant strong deformation retract.*

*Proof.* Let  $h$  be a homeomorphism from  $G \backslash X$  to  $D \subset \mathbb{R}^n$  such that the images of the strata of  $G \backslash X$  under  $h$  are convex. Define  $H' : \mathbb{R}^n \times \mathbb{R}^n \times I \rightarrow \mathbb{R}^n \times \mathbb{R}^n$  by  $H'((a, b), t) = (a, ta + (1 - t)b)$ . Notice that  $H'(a, a, t) = (a, a)$  for every  $a \in \mathbb{R}^n$  and every  $t \in I$ . Let  $S = \{(\sigma(0), \sigma(1)) \mid \sigma \in P_{\text{sp}}(G \backslash X)\}$ , and let  $H = (h \times h)^{-1} \circ H' \circ ((h \times h)|_S \times \text{id}_I)$ . Since the images of the strata of  $G \backslash X$  under  $h$  are convex,  $H : S \times I \rightarrow S$  is a homotopy such that  $H_0 \circ (\rho \times \rho)|_{\tilde{S}} = (\rho \times \rho)|_{\tilde{S}} \circ \text{id}_{\tilde{S}}$  and  $H((\rho \times \rho)(\tilde{S}_{(K \times K^g)}) \times I) \subset (\rho \times \rho)(\tilde{S}_{(K \times K^g)})$  for every finite subgroup  $K$  of  $G$  and every  $g \in G$ . Observe that if  $(x, y) \in \tilde{S}$ , then  $(G \times G)_{(x, y)} = G_x \times G_y = K \times K^g$  for some finite subgroup  $K$  of  $G$  and some  $g \in G$ . Therefore, by the Covering Homotopy Theorem (Theorem 4.7), there exists a  $G \times G$ -isovariant homotopy  $\tilde{H} : \tilde{S} \times I \rightarrow \tilde{S}$  covering  $H$  such that  $\tilde{H}_0 = \text{id}_{\tilde{S}}$ . Since  $(\rho \times \rho)^{-1}(\Delta(G \backslash X)) = F(X)$  it follows that  $\tilde{H}_1(\tilde{S}) \subset F(X)$ . Thus,  $\tilde{H}$  is the desired homotopy.  $\square$

**Corollary 5.2.** *Let  $G$  be a discrete group and  $X$  a proper  $G$ -space. Assume that  $G \backslash X$  is homeomorphic to a subset of  $\mathbb{R}^n$  for some  $n$ , and that the images of the strata of  $G \backslash X$  in  $\mathbb{R}^n$  are convex. If  $F(X)$  is path connected, then  $F(X)_0 = F(X) \hookrightarrow \tilde{S}$  is an isovariant strong deformation retract.*

Next we determine when  $F(X)$  is path connected.

**Theorem 5.3.** *Let  $G$  be a discrete group and  $X$  a path connected  $G$ -space. Then,  $F(X)$  is path connected if every element of  $G$  can be expressed as a product of elements each of which fixes some point in  $X$ . If, in addition,  $G$  acts properly on  $X$ , then the converse is true.*

*Proof.* Let  $S = \{s \in G \mid sy = y \text{ for some } y \in X\}$ . Clearly, if  $s \in S$  and  $y \in X$  such that  $sy = y$ , then  $A_X(s, y) = A_X(1, y)$ . Since  $X$  is path connected, this implies that  $A_X(S \times X) \subset F(X)$  is path connected.

Suppose  $S$  generates  $G$ . Let  $(g, x) \in G \times X$  be given. We will show that there is a path in  $F(X)$  connecting  $A_X(g, x)$  to a point in  $A_X(S \times X)$ . Write  $g = s_n \cdots s_2 s_1$ , where  $s_i \in S$ . For each  $i$ , there is an  $x_i \in X$  such that  $s_i x_i = x_i$ . Therefore,

$$A_X(g, x_1) = A_X(gs_1^{-1}, x_1), \text{ and } A_X(gs_1^{-1} \cdots s_i^{-1}, x_{i+1}) = A_X(gs_1^{-1} \cdots s_{i+1}^{-1}, x_{i+1})$$

for each  $i$ ,  $1 \leq i \leq n-1$ . Since  $X$  is path connected,  $A_X(\{h\} \times X)$  is path connected for every  $h \in G$ . Thus,  $A_X(g, x)$  and  $A_X(1, x_n)$  are connected by a path in  $F(X)$ .

Now assume that  $G$  acts properly on  $X$ , and  $F(X)$  is path connected. Let  $N$  be the subgroup of  $G$  generated by  $S$ . Since  $S$  is closed under conjugation,  $N$  is a normal subgroup of  $G$ . Therefore,  $G/N$  acts on  $N \backslash X$  by  $gN \cdot \rho(x) = \rho(gx)$ , where  $\rho : X \rightarrow N \backslash X$  is the orbit map. It is easy to check that the action is free. The fact that  $G$  acts properly on  $X$  implies that  $N$  acts properly on  $X$  and that  $X$  is Hausdorff; furthermore,  $N \backslash X$  is Hausdorff [4, III, 4.2, Proposition 3]. Recall that a discrete group  $G$  acts properly on a Hausdorff space  $X$  if and only if for every pair of points  $x, y \in X$ , there is a neighborhood  $V_x$  of  $x$  and a neighborhood  $V_y$  of  $y$  such that the set of all  $g \in G$  for which  $gV_x \cap V_y \neq \emptyset$  is finite [4, III, 4.4, Proposition 7]. This implies that  $G/N$  acts properly on  $N \backslash X$ . Therefore,  $A_{G/N} : G/N \times N \backslash X \rightarrow N \backslash X \times N \backslash X$  is a homeomorphism onto its image  $F(N \backslash X)$ . Thus,  $F(N \backslash X)$  is path connected if and only if  $G/N$  is trivial. Since the map  $\rho_F : F(X) \rightarrow F(N \backslash X)$ , defined by  $\rho_F(x, gx) = (\rho(x), \rho(gx))$ , is onto and  $F(X)$  is path connected, it follows that  $G/N$  is trivial. That is,  $G = N$ .  $\square$

An immediate consequence of this theorem is the following.

**Corollary 5.4.** *Let  $G$  be a discrete group and  $\mathcal{F}$  a family of subgroups of  $G$ . If there exists a set of generators,  $S$  of  $G$ , with the property that for every  $s \in S$ , there is an  $H \in \mathcal{F}$  such that  $s \in H$ , then  $F(E_{\mathcal{F}}G)$  is path connected.*

**Example 5.5** (The infinite dihedral group). Let  $G = D_{\infty} = \langle a, b \mid a^2 = 1, aba^{-1} = b^{-1} \rangle$  and  $X = \mathbb{R}$ , where  $a$  acts by reflection through zero and  $b$  acts by translation by 1. Since  $\mathbb{R}$  is a model for  $\underline{E}D_{\infty}$  and  $D_{\infty}$  is generated by two elements of order two, namely  $a$  and  $ab$ ,  $F(\mathbb{R})$  is path connected by Corollary 5.4. The quotient of  $\mathbb{R}$  by  $D_{\infty}$  is homeomorphic to the closed interval  $[0, \frac{1}{2}]$ . The strata are  $\{0\}$ ,  $\{\frac{1}{2}\}$  and  $(0, \frac{1}{2})$ . Therefore, Corollary 5.2 implies that  $F_0(\mathbb{R}) \hookrightarrow \tilde{S}$  is an isovariant strong deformation retract.

**Example 5.6** (Triangle groups). Let

$$G = \langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^p = (bc)^q = (ca)^r = 1 \rangle,$$

where  $p, q, r$  are natural numbers such that  $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} \leq 1$ . The group  $G$  can be realized as a group of reflections through the sides of a euclidean or hyperbolic triangle whose interior angles measure  $\frac{\pi}{p}$ ,  $\frac{\pi}{q}$  and  $\frac{\pi}{r}$ , where the generators  $a$ ,  $b$  and  $c$  act by reflections

through the corresponding sides. Thus, the triangle group  $G$  produces a tessellation of the euclidean or hyperbolic plane by these triangles. Therefore, this plane is a model for  $\underline{EG}$ , whose quotient,  $D$ , is equivalent to the given triangle. By Corollary 5.4,  $F(\underline{EG})$  is path connected. There are seven strata of  $D$ , namely  $\mathring{D}$ ,  $\mathring{S}_a$ ,  $\mathring{S}_b$ ,  $\mathring{S}_c$ , and each of the three vertices, where  $\mathring{D}$  denotes the interior of  $D$ , and  $\mathring{S}_a$ ,  $\mathring{S}_b$ , and  $\mathring{S}_c$  are the interiors of the sides of the triangle,  $S_a$ ,  $S_b$ , and  $S_c$ , respectively, through which  $a$ ,  $b$ , and  $c$  reflect. It follows from Corollary 5.2 that  $F_0(\underline{EG}) \hookrightarrow \tilde{S}$  is an isovariant strong deformation retract.

*Remark 5.7.* Let  $X$  be a  $G$ -space and  $Y$  an  $H$ -space. Clearly,  $F_{G \times H}(X \times Y) \cong F_G(X) \times F_H(Y)$ , and  $F_{G \times H}(X \times Y)_0 \cong F_G(X)_0 \times F_H(Y)_0$ . (Here, the group that is acting has been added to the notation of the configuration space.) Furthermore, since  $(x, y)$  and  $(x', y')$  are in the same stratum of  $X \times Y$  if and only if  $x$  and  $x'$  are in the same stratum of  $X$  and  $y$  and  $y'$  are in the same stratum of  $Y$ , it follows that  $\tilde{S}_{X \times Y} \cong \tilde{S}_X \times \tilde{S}_Y$ . Therefore, if  $F_G(X)_0 \hookrightarrow \tilde{S}_X$  is a  $G$ -isovariant strong deformation retraction and  $F_H(Y)_0 \hookrightarrow \tilde{S}_Y$  is an  $H$ -isovariant strong deformation retraction, then  $F_{G \times H}(X \times Y)_0 \hookrightarrow \tilde{S}_{X \times Y}$  is a  $G \times H$ -isovariant strong deformation retraction. This observation produces interesting examples for which Theorem 4.26 is true. If  $X = \mathbb{R}$ ,  $G = \mathbb{Z}$ ,  $Y = \mathbb{R}$  and  $H = D_\infty$ , then  $F_{\mathbb{Z} \times D_\infty}(\mathbb{R} \times \mathbb{R}) \cong F_{\mathbb{Z}}(\mathbb{R}) \times F_{D_\infty}(\mathbb{R})$  is not path connected, since  $F_{\mathbb{Z}}(\mathbb{R})$  is not path connected. Moreover,  $F_{\mathbb{Z} \times D_\infty}(\mathbb{R} \times \mathbb{R})_0 \neq \Delta(\mathbb{R})$  and  $F_{\mathbb{Z} \times D_\infty}(\mathbb{R} \times \mathbb{R})_0 \neq F_{\mathbb{Z} \times D_\infty}(\mathbb{R} \times \mathbb{R})$ . Despite this, Theorem 4.26 applies to  $\mathbb{Z} \times D_\infty$ .

## 6. A COMPARISON OF $G \backslash F(EG)$ AND $G \backslash F(\underline{EG})$

In this section we examine the map  $\mathfrak{N}(G, \{1\}) \rightarrow \mathfrak{N}(G, \mathcal{F})$ , where  $G$  is a discrete group and  $\mathcal{F}$  is the family of finite subgroups of  $G$ . This enables us to compute the induced map  $HH_*(\mathbb{Z}G) \rightarrow HH_*^{\mathcal{F}}(\mathbb{Z}G)$ .

Let  $E$  be a model for  $EG$  and  $\underline{E}$  be a model for the universal space for proper  $G$ -actions. Then,  $G \backslash F(E)$  is homeomorphic to  $\mathfrak{N}(G, \{1\})$ , and  $\mathfrak{N}(G, \mathcal{F})$  is homeomorphic to  $G \backslash F(\underline{E})$  by Corollary 3.7. The universal property of  $\underline{E}$  implies that there is a  $G$ -equivariant map,  $f : E \rightarrow \underline{E}$ , that is unique up to  $G$ -homotopy equivalence. Then  $F(f) : F(E) \rightarrow F(\underline{E})$  induces a map  $\bar{f} : G \backslash F(E) \rightarrow G \backslash F(\underline{E})$ . Note that for a different choice of  $f$ , the induced map will be homotopy equivalent to  $\bar{f}$ . The corresponding map on homology groups is denoted  $\bar{f}_* : HH_*(\mathbb{Z}G) \rightarrow HH_*^{\mathcal{F}}(\mathbb{Z}G)$ . Recall that

$$\bar{A}_E : G \backslash (G \times E) \rightarrow G \backslash F(E)$$

is a homeomorphism, since  $G$  acts freely and properly on  $E$  (Proposition 3.10). By Proposition 3.11, there is a homeomorphism

$$h : \coprod_{C(g) \in \text{conj}(G)} Z(g) \backslash E \rightarrow G \backslash (G \times E),$$

which sends the orbit  $Z(g) \cdot x$  to the orbit  $G \cdot (g, x)$ . This produces a map

$$\phi : \coprod_{C(g) \in \text{conj}(G)} Z(g) \backslash E \rightarrow G \backslash F(\underline{E}),$$

where  $\phi = \bar{f} \circ \bar{A}_E \circ h$ . That is, the image of  $Z(g) \cdot x$  under  $\phi$  is  $G \cdot (f(x), g \cdot f(x))$ , where  $g$  is in  $G$  and  $x$  is in  $E$ . Thus, we have the following commutative diagram.

$$\begin{array}{ccc} HH_*(\mathbb{Z}G) & \xrightarrow{\bar{f}_*} & HH_*^{\mathcal{F}}(\mathbb{Z}G) \\ \uparrow \cong & \nearrow \phi_* & \\ \bigoplus_{C(g) \in \text{conj}(G)} H_*(BZ(g); \mathbb{Z}) & & \end{array}$$

If  $H$  is a finite group, then the Sullivan Conjecture, proved by Miller [17], implies that a map from  $BH$  to a finite dimensional CW complex is null homotopic. If  $\underline{E}$  is finite dimensional, then  $F(\underline{E})$  is homotopy equivalent to a finite dimensional CW complex. Thus, if  $Z(g)$  is finite, then the image of  $H_*((BZ(g); \mathbb{Z}))$  under  $\phi_*$  is zero.


For an illustrative example, consider the infinite dihedral group,  $D_\infty = \langle a, b \mid a^2 = 1, aba^{-1} = b^{-1} \rangle$ . Let  $\underline{E} = \mathbb{R}$ , where  $a$  acts by reflection through zero and  $b$  acts by translation by 1. That is,  $ax = -x$  and  $bx = x + 1$ . The space  $F(\mathbb{R})$  is the image of  $A_\mathbb{R} : D_\infty \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ . Thus,  $F(\mathbb{R}) = \{(x, gx) \mid x \in \mathbb{R} \text{ and } g \in D_\infty\}$ . Every element of  $D_\infty$  can be expressed as  $b^j$  or  $ab^j$ , for some  $j$  in  $\mathbb{Z}$ . Since  $b^j x = x + j$  and  $ab^j x = -x - j$ ,  $F(\mathbb{R}) \subset \mathbb{R}^2$  is the union of the lines of slope 1 and -1 that cross the  $y$ -axis at an integer. A picture of  $D_\infty \backslash F(\mathbb{R})$  is given in Figure (1) below.

To see that this is in fact the picture, consider the diagonal action of  $\langle b \rangle$  on  $\mathbb{R}^2$ . The orbit of the set

$$D = \{(x, y) \mid x \in \mathbb{R} \text{ and } -x - 1 \leq y \leq -x + 1\}$$

under this action is all of  $\mathbb{R}^2$ . Observe that the lines  $y = -x - 1$  and  $y = -x + 1$  get identified in the quotient of  $\mathbb{R}^2$  by  $\langle b \rangle$  and that the rest of the set is mapped injectively into the quotient. Thus,  $\langle b \rangle \backslash \mathbb{R}^2$  is an infinite cylinder. Since  $a$  acts on the set  $D$  by a rotation of  $180^\circ$ , we see that the quotient  $D_\infty \backslash \mathbb{R}^2 = \langle a \rangle \backslash (\langle b \rangle \backslash \mathbb{R}^2)$  is obtained from



FIGURE 1. The space  $D_\infty \setminus F(\mathbb{R})$ .

$\{(x, y) \in D \mid y \geq x\}$  by identifying the endpoints of the line segments  $y = x + t$ , where  $t \geq 0$  (that is, the points  $(\frac{-t-1}{2}, \frac{t-1}{2})$  and  $(\frac{-t+1}{2}, \frac{t+1}{2})$ ), as well as by identifying the points  $(x, x)$  and  $(-x, -x)$ , where  $-1/2 \leq x \leq 1/2$ . Thus,  $D_\infty \setminus \mathbb{R}^2$  looks like an “infinite chisel”, and  $D_\infty \setminus F(\mathbb{R}) \subset D_\infty \setminus \mathbb{R}^2$  is as shown above.

The non-trivial finite subgroups of  $D_\infty$  are of the form  $\langle ab^i \rangle$ , where  $i \in \mathbb{Z}$ . For each  $i$ ,  $\langle ab^i \rangle$  fixes  $-i/2 \in \mathbb{R}$ . Beginning with the action of  $D_\infty$  on  $\mathbb{R}$ , construct a model for  $ED_\infty$  by replacing each half-integer with an  $S^\infty$ . Denote this “string of pearls” model for  $ED_\infty$  by  $E$ , and let  $f : E \rightarrow \underline{E}$  be the equivariant map that collapses each  $S^\infty$  to a point. The conjugacy classes of  $D_\infty$  are:

$$\begin{aligned} C(1) &= \{1\} \\ C(a) &= \{ab^{2i} : i \in \mathbb{Z}\} \\ C(ab) &= \{ab^{2i+1} : i \in \mathbb{Z}\} \\ C(b^j) &= \{b^j, b^{-j}\}, \quad j \in \mathbb{N}. \end{aligned}$$

The corresponding centralizers are

$$\begin{aligned} Z(1) &= D_\infty \\ Z(a) &= \{1, a\} \\ Z(ab) &= \{1, ab\} \\ Z(b^j) &= \langle b \rangle, \quad j \in \mathbb{N}. \end{aligned}$$

Note that  $D_\infty \setminus E$  is an “interval” with an  $\mathbb{R}P^\infty$  at each end;  $\langle a \rangle \setminus E$  is a “ray” that begins with an  $\mathbb{R}P^\infty$  at 0 and has an  $S^\infty$  at every positive half-integer;  $\langle ab \rangle \setminus E$  is a “ray” that begins with an  $\mathbb{R}P^\infty$  at  $1/2$  and has an  $S^\infty$  at every other positive half-integer; and  $\mathbb{Z} \setminus E$  is a “circle” with two  $S^\infty$ ’s in place of vertices.

The image of  $\phi$  is broken into the pieces

$$(4) \quad \phi(D_\infty \cdot x) = D_\infty \cdot (f(x), f(x))$$

$$(5) \quad \phi(Z(a) \cdot x) = D_\infty \cdot (f(x), -f(x))$$

$$(6) \quad \phi(Z(ab) \cdot x) = D_\infty \cdot (f(x), -f(x) - 1)$$

$$(7) \quad \phi(Z(b^j) \cdot x) = D_\infty \cdot (f(x), f(x) + j),$$

where  $j$  is a positive integer and  $x \in E$ . Referring to diagram (1), the base of  $D_\infty \setminus F(E)$  is (4), the pieces (5) and (6) are the sides of  $D_\infty \setminus F(E)$ , and (7) provides each of the circles. Therefore,  $\phi$  is a gluing of the disjoint pieces,  $Z(g) \setminus E$ , after each  $S^\infty$  and each  $\mathbb{R}P^\infty$ , is collapsed to a point. Observe that,

$$HH_*(\mathbb{Z}D_\infty) \cong H_*(BD_\infty; \mathbb{Z}) \oplus H_*(BZ(a); \mathbb{Z}) \oplus H_*(BZ(ab); \mathbb{Z}) \oplus \bigoplus_{j>0} H_*(BZ(b^j); \mathbb{Z}).$$

Since  $Z(a) \cong \mathbb{Z}/2 \cong Z(ab)$ , the Sullivan Conjecture implies that the image of  $H_*(BZ(a); \mathbb{Z})$  and  $H_*(BZ(ab); \mathbb{Z})$  under  $\phi_*$  is zero. By the above analysis,  $\phi(BD_\infty) = D_\infty \setminus \mathbb{R} \cong [0, 1]$ . Therefore, the image of  $H_i(BD_\infty; \mathbb{Z})$  under  $\phi_i$  is 0, for  $i \geq 1$ . The rest of  $HH_i(\mathbb{Z}D_\infty)$  is mapped injectively into  $HH_i^{\mathcal{F}}(\mathbb{Z}D_\infty)$ ,  $i \geq 1$ .

Classical Hochschild homology has been used to study the  $K$ -theory of groups rings via the *Dennis trace*,  $\text{dtr} : K_*(RG) \rightarrow HH_*(RG)$ . In [15], Lück and Reich were able to determine how much of  $K_*(\mathbb{Z}G)$  is detected by the Dennis trace. A natural question is to determine the composition of the Dennis trace with the map  $\bar{f}_* : HH_*(\mathbb{Z}G) \rightarrow HH_*^{\mathcal{F}}(\mathbb{Z}G)$ . From Lück and Reich, we have the following commutative diagram

$$\begin{array}{ccc} H_*^G(\underline{E}; \mathbf{K}_{\mathbb{Z}}) & \xrightarrow{A} & K_*(\mathbb{Z}G) \\ \downarrow & & \downarrow \text{dtr} \\ H_*^G(\underline{E}; \mathbf{HH}_{\mathbb{Z}}) & \xrightarrow{B} & HH_*(\mathbb{Z}G) \end{array}$$

[15, p.595], where the maps  $A$  and  $B$  are *assembly maps* in the equivariant homology theories with coefficients in the connective algebraic  $K$ -theory spectrum,  $\mathbf{K}_{\mathbb{Z}}$ , associated to  $\mathbb{Z}$ , and the Hochschild homology spectrum  $\mathbf{HH}_{\mathbb{Z}}$ , respectively. Each assembly map is

induced by the collapse map  $\underline{E} \rightarrow \text{pt.}$  Lück and Reich use the composition of the Dennis trace with the assembly map in algebraic  $K$ -theory,  $\text{dtr} \circ A$ , to achieve their detection results. In particular, they observe that the assembly map in Hochschild homology factors as

$$\begin{array}{ccc}
 H_*^G(\underline{E}G; \mathbf{HH}_Z) & \xrightarrow{B} & HH_*(ZG) \\
 \uparrow \cong & & \uparrow \cong \\
 \bigoplus_{\substack{C(g) \in \text{conj}(G) \\ \langle g \rangle \in \mathcal{F}}} H_*(BZ(g); \mathbb{Z}) & \hookrightarrow & \bigoplus_{C(g) \in \text{conj}(G)} H_*(BZ(g); \mathbb{Z})
 \end{array}$$

[15, p. 630]. Given the discussion above, in the case  $G = D_\infty$ ,

$$H_*^G(\underline{E}D_\infty; \mathbf{HH}_Z) \cong H_*(BD_\infty; \mathbb{Z}) \oplus H_*(BZ(a); \mathbb{Z}) \oplus H_*(BZ(ab); \mathbb{Z}).$$

Therefore,  $\bar{f}_* \circ B = 0$ , which implies that the image of  $\bar{f}_* \circ \text{dtr} \circ A$  is zero.

We conclude with speculation about a possible geometric application of the groups  $HH_*^{\mathcal{F}}(\mathbb{Z}G)$ . Associated to a parametrized family of self-maps of a manifold  $M$ , there are geometrically defined “intersection invariants,” in particular, the framed bordism invariants of Hatcher and Quinn [10], which take values in abelian groups that are known to be related to the Hochschild homology groups  $HH_*(\mathbb{Z}G)$ , where  $G$  is the fundamental group of  $M$  [9]. It appears plausible that the groups  $HH_*^{\mathcal{F}}(\mathbb{Z}G)$ , where  $\mathcal{F}$  is the family of finite subgroups, could play an analogous role in the yet to be developed homotopical intersection theory of orbifolds.

## REFERENCES

- [1] R. Ayala, F.F. Lasheras, A. Quintero, *The equivariant category of proper  $G$ -spaces*, Rocky Mountain J. Math. **31** (2001), 1111–1132.
- [2] H. Biller, *Characterizations of proper actions*, Math. Proc. Camb. Phil. Soc. **136** (2004), 429–439.
- [3] C.J.R. Borges, *On stratifiable spaces*, Pacific J. Math. **17** (1966), 1–16.
- [4] N. Bourbaki, *Elements of Mathematics: General Topology, Part I* (English edition), Addison Wesley, Reading MA, 1966.
- [5] G.E. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, New York, 1972.
- [6] R. Cauty, *Sur les espaces d'applications dans les CW-complexes*, Arch. Math. (Basel) **27** (1976), 306–311.
- [7] J. Davis and W. Lück, *Spaces over a category and assembly maps in isomorphism conjectures in  $K$ - and  $L$ -theory*,  $K$ -theory **15** (1998), 201–252.
- [8] J.J. Duistermaat and J.A.C. Kolk, *Lie groups*, Springer-Verlag, Berlin, 2000.
- [9] R. Geoghegan and A. Nicas, *Parametrized Lefschetz-Nielsen fixed point theory and Hochschild homology traces*, Amer. J. Math. **116** (1994), 397–446.

- [10] A. Hatcher and F. Quinn, *Bordism invariants of intersections of submanifolds*, Trans. Amer. Math. Soc. **200** (1974), 327–344.
- [11] B. Hughes, *Stratified path spaces and fibrations*, Proc. Roy. Soc. Edinburgh **129A** (1999), 351–384.
- [12] J.-L. Loday, *Cyclic homology*, second ed., Springer-Verlag, New York, NY, 1998.
- [13] W. Lück, *Transformation Groups and Algebraic K-Theory*, Lecture Notes in Mathematics vol. 1408, Springer-Verlag, Berlin, 1989.
- [14] W. Lück, *Survey on classifying spaces for families of subgroups*. In Infinite groups: geometric, combinatorial and dynamical aspects, vol. 248 of Progr. Math., Birkhäuser, Basel, 2005, pp. 269–322.
- [15] W. Lück and H. Reich, *Detecting K-theory by cyclic homology*, Proc. London Math. Soc. (3) **93** (2006), 593–634.
- [16] A.T. Lundell and S. Weingram, *The Topology of CW Complexes*, Van Nostrand Reinhold Co., New York, 1969.
- [17] H. Miller, *The Sullivan conjecture on maps from classifying spaces*, Ann. of Math. **120** (1984), 39–87.
- [18] J. Milnor, *Construction of universal bundles, I*, Ann. of Math. **63** (1956), 272–284.
- [19] R.S. Palais, *The classification of G-spaces*, Mem. Amer. Math. Soc., No. 36, 1960.
- [20] R.S. Palais, *On the existence of slices for actions of non-compact Lie groups*, Ann. of Math. **73** (1961), 295–323.
- [21] E.H. Spanier, *Algebraic Topology*, McGraw-Hill, New York, 1966.
- [22] R. Talbert, *An isomorphism between Bredon and Quinn homology*, Forum Math. **11** (1999), 591–616.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO, CANADA L8S 4K1

*E-mail address:* `nicas@mcmaster.ca`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHNS UNIVERSITY, 8000 UTOPIA PKWY, JAMAICA, NY 11439, USA

*E-mail address:* `rosenthd@stjohns.edu`